arXiv:1205.5544v3 [math.SG] 8 Oct 2012

LEGENDRIAN AMBIENT SURGERY AND LEGENDRIAN CONTACT HOMOLOGY

GEORGIOS DIMITROGLOU RIZELL

Abstract. Let L be a Legendrian submanifold of a contact manifold Y and let $S \subset L$ be a framed sphere bounding a subcritical isotropic disk in $Y \setminus L$. We may perform an ambient surgery on L along S, obtaining a Legendrian submanifold $L_S \subset Y$ which is Lagrangian cobordant to L. We produce an isomorphism of the Legendrian contact homology algebra of L_S with an algebra obtained from the algebra of L after twisting the differential by a count of holomorphic disks with boundary points mapping to S. This isomorphism induces a one-to-one correspondence between the augmentations for the algebras of L and L_S.

1. Introduction

A contact manifold is a $(2n+1)$-dimensional manifold Y together with a maximally non-integrable hyper-plane field ξ. For us, this hyper-plane field will be given by

$$\xi = \ker(\lambda),$$

where λ is a fixed global 1-form called the *contact form*, which consequently must satisfy the property that $\lambda \wedge (d\lambda)^n$ is everywhere non-zero.

The *Reeb vector field* on (Y, λ) is the vector field R defined by the requirements

$$\iota_R d\lambda = 0, \;\; \lambda(R) = 1.$$

The definition of R clearly depends on the choice of λ.

We have to make some technical assumptions on Y in order for the results in [Ekh2] to hold. We refer to Section 2 for more details. In particular, the conditions are satisfied for the *jet-space* of a smooth manifold M. This is the contact manifold

$$(J^1(M) \simeq T^*M \times \mathbb{R}, dz + \theta),$$

where z is the coordinate on the $\mathbb{R}$-factor and where θ is the Liouville form on T^*M.

An embedded submanifold $L \subset Y$ is called *isotropic* if $TL \subset \xi$, which implies that $\dim L \leq n$. In the case $\dim L = n$ we say that L is *Legendrian*, and in the case $\dim L < n$ we say that L is *subcritical isotropic*. A *Reeb chord* on L is an integral curve of R starting and ending on two different sheets of L. Unless stated otherwise, we will make the assumption that a Legendrian submanifold $L \subset Y$ is closed.

Let (Y, λ) be a contact manifold. The exact symplectic manifold

$$(\mathbb{R} \times Y, d(e^t \lambda)),$$

where t is a coordinate on the $\mathbb{R}$-factor, is called the *symplectization* of (Y, λ). By an exact Lagrangian cobordism from the Legendrian manifold $L_1 \subset Y$ to $L_2 \subset Y$, we mean an exact Lagrangian submanifold $V \subset \mathbb{R} \times Y$ (that is, $e^t \lambda$ is exact when pulled back to V), which coincides with

$$((-\infty, -N) \times L_1) \cup ((N, +\infty) \times L_2), \;\; N \gg 0$$

outside of a compact subset.

For a Legendrian submanifold $L \subset Y$, the trivial cylinder $\mathbb{R} \times L \subset \mathbb{R} \times Y$ is an exact Lagrangian cobordism from L to L. Observe that the relation of being exact Lagrangian cobordant is not a symmetric relation.

Recall that a symplectic manifold (X, ω) has many *compatible* almost complex structures, that is, $J \in \mathrm{End}(TX)$ such that $J^2 = -Id$ and $\omega(\cdot, J\cdot)$ is a Riemannian metric. An almost complex structure J on $\mathbb{R} \times Y$ is called *cylindrical* if it is compatible with $d(e^t \lambda)$, translation invariant under the natural $\mathbb{R}$-action on $\mathbb{R} \times Y$, leaves the contact planes invariant, and satisfies $J\partial_t = R$, where $R \in TY$ is the Reeb vector field with respect to λ.

Symplectic Field Theory, introduced in [EGH] by Eliashberg, Givental and Hofer, provides a framework for algebraic invariants, for contact manifolds and Legendrian submanifolds (among others), which encode information about related spaces of pseudo-holomorphic curves.

A *Legendrian isotopy* is a smooth 1-parameter family L_t of Legendrian embeddings. Legendrian Contact homology is a Legendrian isotopy invariant obtained from Symplectic Field Theory, which was defined in [EGH] by Eliashberg, Givental and Hofer, and in [Che] by Chekanov.

In Section 2 we give an outline of the definition of Legendrian contact homology. Roughly speaking, it is constructed as follows. For a compact Legendrian submanifold $L \subset Y$ and a cylindrical almost complex structure J on $\mathbb{R} \times Y$, one defines its contact homology algebra

$$(\mathcal{A}(L), \partial)$$

which is a differential graded algebra (DGA for short). The algebra $\mathcal{A}(L)$ is unital, non-commutative, and freely generated by the Reeb chords on L. The differential counts rigid (up to translation) J-holomorphic disks in $\mathbb{R} \times Y$ with boundary on $\mathbb{R} \times L$, one boundary puncture asymptotic to a Reeb chords on L at $+\infty$ and several boundary punctures asymptotic to Reeb chords on L at $-\infty$. The homotopy type of this DGA is invariant under Legendrian isotopy as well as the choice of a generic cylindrical almost complex structure.

By a standard construction in Symplectic Field Theory, an exact Lagrangian cobordism V from L_1 to L_2 induces a chain map

$$\Phi \colon\; (\mathcal{A}(L_2), \partial) \to (\mathcal{A}(L_1), \partial)$$

which is defined by counting rigid J-holomorphic disks in $\mathbb{R} \times Y$ having boundary on V, one boundary puncture asymptotic to a Reeb chord on L_2 at $+\infty$, and several boundary punctures asymptotic to Reeb chords on L_1 at $-\infty$.

In Section 3 we describe the following construction. Let $L \subset Y$ be a Legendrian submanifold and $S \subset L$ a framed embedded k-sphere, where $k < n$, such that S bounds an isotropic $(k+1)$-disk $D \subset Y \setminus L$. Let L_S be the smooth manifold obtained from L by surgery along S. Under some additional hypotheses on D which are discussed in Section 3, one can perform a *Legendrian ambient surgery* on L along S, resulting in a Legendrian embedding of L_S into an arbitrarily small neighborhood of $L \cup D$. The Reeb chords on L_S will consist of Reeb chords coinciding with the Reeb chords on L, together with a new Reeb chord c_{new}, of arbitrary small action, situated in the middle of the newly added handle.

The construction moreover provides an exact Lagrangian cobordism $V_\epsilon \subset \mathbb{R} \times Y$ from L to L_S. This cobordism is diffeomorphic to the manifold obtained by a $(k+1)$-handle attachment on $(-\infty, -1] \times L$ along

$$S \subset L = \partial((-\infty, -1] \times L)$$

with the prescribed framing. Observe that the symplectization coordinate restricted to V_ϵ has a unique critical point of index $k+1$.

Let

$$\Phi \colon (\mathcal{A}(L_S), \partial) \to (\mathcal{A}(L), \partial)$$

be the morphism of DGAs induced by V_ϵ. Fist, observe that the disk-counts in the proofs of Theorem 1.4 and Proposition 7.3 give the following result.

Proposition 1.1. *There is a cylindrical almost complex structure on $\mathbb{R} \times Y$ for which Φ is a surjection having kernel $\langle c_{new} \rangle$ in the case $k < n-1$, and $\langle c_{new} - 1 \rangle$ in the case $k = n-1$.*

The goal of this paper is to compute the DGA $(\mathcal{A}(L_S), \partial)$ in terms of data on L, as well as the morphism Φ in terms of this identification.

In the following, we make the additional assumption that $k < n-1$. In Section 4, given k-sphere $S \subset L$ with framed normal bundle N, we define the DGA

$$(\mathcal{A}(L, S), \partial_{S,N}),$$

which is supposed to model the DGA of L_S obtained by Legendrian ambient surgery on L along S, in the case when it can be performed.

The algebra $\mathcal{A}(L, S)$ is unital, non-commutative and freely generated over $\mathbb{Z}_2$ by the Reeb chords on L together with a formal generator s in degree $n-k-1$. The formal generator should be thought of as corresponding to the Reeb chord c_{new} appearing after a Legendrian ambient surgery along S.

The differential, which depends on the choice of framing for N, can roughly be described as follows. We fix a non-vanishing normal vector-field $\mathbf{v}$ to S which is homotopic to a constant vector-field with respect to the chosen framing. We set $\partial_{S,N}(s) = 0$. For a Reeb chord generator a we define $\partial_{S,N}(a)$ to have coefficient in front of the word

$$s^{w_1} b_1 s^{w_2} \cdot \ldots \cdot s^{w_m} b_{m+1} s^{w_{m+1}}$$

given by the number of points, counted modulo 2, of the union of all 0-dimensional spaces $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v})/\mathbb{R}$ for different $A \in H_1(L)$ and $\mathbf{w} \in (\mathbb{Z}_{\geq 0})^{m+1}$. The moduli space $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v})/\mathbb{R}$, defined in Section 8, consists of the J-holomorphic disks in the symplectization $\mathbb{R} \times Y$, defined up to translation, having:

- boundary mapping to $\mathbb{R} \times L$ in homology class $A \in H_1(L)$,
- one boundary puncture p_0 asymptotic to the Reeb chord a at $+\infty$,
- m ordered boundary punctures $p_1, \ldots, p_m$ (where the order is determined by the orientation of $\partial D \setminus \{p_0\}$), which are asymptotic to the Reeb chords $b_1, \ldots, b_m$, respectively, at $-\infty$, and
- w_i points on the boundary arc between the punctures p_i and p_{i+1} mapping to $\mathbb{R} \times S$ (where we set $p_{m+1} := p_0$). We count each point with multiplicity $l+1$, where l is the order of tangency of the boundary to $\exp_{\mathbb{R}\times S}(\mathbb{R}\mathbf{v})$ in case when the tangent of the boundary has a positive component in the $\mathbf{v}$-direction, and where otherwise $l = 0$.

Theorem 1.2. *Let $S \subset L$ be a k-dimensional sphere with trivial normal bundle, such that $k < n-1$. There are cylindrical almost complex structures on $\mathbb{R} \times Y$ for which*

$$(\mathcal{A}(L, S), \partial_{S,N})$$

is a well-defined DGA. Moreover,

$$(\mathcal{A}(L), \partial) \simeq (\mathcal{A}(L, S)/\langle s \rangle, \partial_{S,N}).$$

When counting disks with multiple boundary-point constraints on $\mathbb{R} \times S$, the remark below shows that we also need to consider pseudo-holomorphic disks with higher order jet-constraints at the boundary. We set up the relevant moduli spaces in Section 8. Since we are dealing with jet-constraints of pseudo-holomorphic disks, we will use almost complex structures which are integrable in a neighborhood of $\mathbb{R} \times S$ to be able to locally model standard solutions. An alternative approach for dealing with jet-constraints on boundary points is conducted in [Zeh].

We also relate the moduli spaces of disks with jet-constraints on $\mathbb{R} \times S$ to the moduli spaces defined in Section 9, which consist of disks with boundary-point constraints on the parallel copies $\exp_{\mathbb{R}\times S}(i\delta\mathbf{v})$ of $\mathbb{R} \times S$, where $i \in \mathbb{N}$ and $\delta > 0$, and where exp denotes the exponential map on $\mathbb{R} \times L$ for some choice of metric. These spaces are in general easier to define, especially when the expected dimension is positive.

Remark 1.3. Consider a count of rigid solutions of J-holomorphic disks with the constraint that d geometric boundary points map to $\mathbb{R} \times S$. One can consider the following perturbation of the problem. For each $\delta > 0$, we require that d points map to the d parallel copies $\mathbb{R} \times S_i := \exp_{\mathbb{R}\times S}(i\delta\mathbf{v})$,

$i = 1, \dots, d$, where $\mathbf{v}$ is a normal vector-field to S and exp denotes the exponential map on $\mathbb{R} \times L$ for some choice of metric, such that the i:th boundary point is mapped to $\mathbb{R} \times S_i$.

For small enough $\delta > 0$, this perturbation of the problem should have the same number of solutions as the original one. However, when we recover the original problem by letting $\delta \to 0$, there may be a sequence of solutions for which $l+1 > 1$ consecutive boundary points mapping to $\mathbb{R} \times S_i, \dots, \mathbb{R} \times S_{i+l}$ converge to a single boundary point in the limit. Consequently, the solution in the limit has a tangency to $\exp_{\mathbb{R}\times S}(\mathbb{R}\mathbf{v})$ of order at least $l > 0$ at this boundary point, but has less than d geometric boundary points mapping to $\mathbb{R} \times S$. From this example we conclude that we have to keep track of the order of tangency to $\exp_{\mathbb{R}\times S}(\mathbb{R}\mathbf{v})$ in the original problem.

In Section 5, we show the following connection between the above DGA and $(\mathcal{A}(L_S), \partial)$.

Theorem 1.4. *Suppose that $k < n-1$ and that L_S is obtained from L by a Legendrian ambient surgery along S. Furthermore, suppose that L and L_S have finitely many generic Reeb chords. For some cylindrical almost complex structure J on $\mathbb{R} \times Y$ there is a tame isomorphism of DGAs*

$$\Psi \colon (\mathcal{A}(L_S), \partial) \to (\mathcal{A}(L, S), \partial_{S,N}),$$

such that the DGA morphism

$$\Phi \colon (\mathcal{A}(L_S), \partial) \to (\mathcal{A}(L), \partial)$$

induced by the exact Lagrangian cobordism V_ϵ corresponds to the natural quotient projection

$$\Phi \circ \Psi^{-1} \colon (\mathcal{A}(L, S), \partial_{S,N}) \to (\mathcal{A}(L, S)/\langle s \rangle, \partial_{S,N}) \simeq (\mathcal{A}(L), \partial).$$

The main part of the proof is the construction of the map Ψ. It is defined by counting rigid pseudo-holomorphic disks in $\mathbb{R} \times Y$ having boundary on V_ϵ and boundary points evaluating to parallel copies of the core disk $\widetilde{S} \subset V_\epsilon$ of the handle attachment V_ϵ, where $\widetilde{S}$ coincides with $(-\infty, -1) \times S$ outside of a compact set (see Section 5.1).

Finally, we have the following consequence of the above theorem. Let $\mu \in H^1(V_\epsilon; \mathbb{Z})$ be the Maslov class of V_ϵ.

Corollary 1.5. *Under the assumptions of the above theorem, if moreover*

$$(n - k - 1) \notin \mu(H_1(V_\epsilon)),$$

then there is a bijection between the sets of augmentations of $(\mathcal{A}(L), \partial)$ and $(\mathcal{A}(L_S), \partial)$.

Remark 1.6. In the case $Y = J^1(M)$ and $\mu(V_\epsilon) = 0$, under the above assumption it follows that any generating family defined on either L or L_S can be extended over the cobordism V_ϵ. In a forthcoming work by Bourgeois, Sabloff and Traynor, formulas are proven which relate the generating family homologies before and after an ambient Legendrian surgery. Generating family homology is a Legendrian isotopy invariant, introduced in [PC] and [FR]. In [FR] it is proven that generating family homology coincides with linearized Legendrian contact homology in the case $Y = J^1(\mathbb{R})$. In higher dimensions, however, not much is known.

There are analogous results in the case when there are infinitely many Reeb chords on L and L_S. However, in this case one has to perform a direct limit construction. We refer to Section 6 for the results.

Remark 1.7. A particular case of Theorem 1.4 allows us to compute the Legendrian contact homology of the *cusp connected sum* $L_1 \# L_2 \subset J^1(M^n)$ of two Legendrian submanifolds $L_1, L_2 \subset J^1(M^n)$ when $n \geq 2$. This operation was described in [EES2, Section 4.2] and is the special case of a Legendrian ambient 0-surgery on $L_1 \cup L_2$. The computation of the Legendrian contact homology of $L_1 \# L_2$ also follows from the results in [HS].

Remark 1.8. The results in this paper can be seen as the first small steps for obtaining a relative version of the results in [BEE], which explains the relation between the Linearized contact homologies of the ends of a Weinstein cobordism, computed by considering pseudo-holomorphic curves determined by a decomposition of the cobordism into standard handles. Linearized contact homology is an invariant of a contact manifold obtained from Symplectic Field Theory (see [EGH]). Similarly, in the relative case, one would like to understand the relations of the Legendrian contact homologies of the ends of a general exact Lagrangian cobordism inside a general Weinstein cobordism in terms of the handle decompositions. However, observe that it is not even clear that a general exact Lagrangian cobordism can be decomposed into a sequence of standard handle-attachments (that is, handles corresponding to Legendrian ambient surgeries).

We end with some remarks regarding the case $k = n - 1$, which is fundamentally different from the above cases. We obtain some partial results in Proposition 7.3. For the case of an ambient 0-surgery on a Legendrian 1-knot inside $J^1(\mathbb{R})$, we refer to [EHK, Lemma 3.15], which gives a computation of the Legendrian contact homology *before* the surgery in terms of data on the knot *after* the surgery.

Acknowledgements. I would like to thank Tobias Ekholm, my Ph. D. advisor, who suggested me the problem and was very helpful during the writing of this article. I would also like to thank Frédéric Bourgeois for pointing out mistakes and providing valuable comments.

2. Background

Legendrian contact homology, which was developed in [EGH] by Eliashberg, Givental and Hofer, and in [Che] by Chekanov, is a theory that associates a differential graded algebra, or DGA for short, to a Legendrian submanifold of a contact manifold. The homotopy type of this DGA is invariant under Legendrian isotopy.

The technical details for this theory have not been worked out in all situations, but both the construction and the invariance have been worked out in $J^1(\mathbb{R}^1)$ by Chekanov in [Che], and in arbitrary contact manifolds on the form $(P\times\mathbb{R}, dz+\theta)$, where $(P, d\theta)$ is an exact symplectic manifold satisfying a convexity assumption, by Ekholm, Etnyre and Sullivan in [EES3]. The invariance for closed contact manifolds (Y, λ) having the property that all periodic Reeb orbits of (Y, λ) are non-degenerate and have Conley-Zehnder index at least 2 follows from the results in [Ekh2].

From now on we will make the assumption that either (Y, λ) is closed and satisfies the conditions above, or on the form $(P \times \mathbb{R}, dz + \theta)$ where P is convex at infinity. For instance, the standard contact form on $J^1(M)$ for a smooth manifold M, where M is either closed or equal to $\mathbb{R}^n$, is of the latter kind.

For a closed Legendrian submanifold $L \subset Y$ we let $\mathcal{Q}(L)$ denote the set of Reeb-chords on L. Throughout this paper we will assume that L is *chord generic*, that is, the Reeb flow takes the tangent plane of L at the starting point of a Reeb chord to a plane transverse to the tangent plane of L at the end point of the Reeb chord. We define the *action* of a Reeb chord c to be

$$\ell(c) := \int_c \lambda.$$

Chord genericity implies that below any given action the set of Reeb-chords on L is a compact, and hence finite, set.

2.1. **The algebra and the grading.**

We now define the underlying algebra of the DGA associated to L. We let

$$\mathcal{A}(L) = \mathbb{Z}_2\langle \mathcal{Q}(L)\rangle$$

be the unital non-commutative algebra over $\mathbb{Z}_2$ which is freely generated by the Reeb chords $\mathcal{Q}(L)$ on L. We fix a *capping path* γ_c for for each Reeb chord c on L, by which we mean a path on L which starts at the end-point of c, and ends at the starting-point of c. We grade each Reeb-chord generator c by

$$|c| = \nu(\gamma_c) - 1,$$

where $\nu(\gamma_c)$ denotes the Conley-Zehnder index (see [EES1] for a definition).

In the case when $Y = J^1(M)$, the following description of the Conley-Zehnder index in terms of the front projection is given in [EES2]. Let f_s and f_e be real-valued functions defined on an open set of M such that the the z-coordinates of the sheets of L at the starting-point and end-point of the Reeb chord c is given by f_s and f_e, respectively. Let $p \in M$ be the projection to M of the Reeb chord c. The Conley-Zehnder index can be expressed as

$$\nu(\gamma_c) = D(\gamma_c) - U(\gamma_c) + \mathrm{index}_p(f_e - f_s), \tag{2.1}$$

where $D(\gamma_c)$ and $U(\gamma_c)$ denote the number of cusp-edges in the front projection traversed by (a generic perturbation of) γ_c in downward and upward direction, respectively, and where $\text{index}_p(f_e - f_s)$ denotes the Morse index at p.

The *Maslov class* will be denoted by $\mu \in H^1(L;\mathbb{Z})$. Again, we refer to [EES1] for a definition. In the case of $Y = J^1(M)$ it may be computed by the formula

$$\mu([\gamma]) = D(\gamma) - U(\gamma), \tag{2.2}$$

where γ is a generic closed curve, and D and U are as above. When the Maslov class is non-zero, the grading has to be taken in the group $\mathbb{Z}/\mu(H_1(L))$.

2.2. **The moduli spaces and compactness.** The differential is defined by counting certain pseudo-holomorphic disks in $\mathbb{R} \times Y$. Here we give a general description of the moduli spaces which we will make us of in this section. These moduli spaces are special cases of the moduli spaces defined in Section 8 with more care.

Let $V \subset \mathbb{R} \times Y$ be an exact Lagrangian cobordism from the Legendrian submanifold L_1 to L_2. Fix a cylindrical almost complex structure J on $\mathbb{R} \times Y$. Let a be a Reeb chord on L_2, $\mathbf{b} = b_1 \cdot \ldots \cdot b_m$ a word of Reeb chords on L_1, $A \in H_1(V)$ be a homology class, and $\dot{D} = D \setminus \{p_0, \ldots, p_m\}$ be the unit disk with $m+1$ boundary points removed. We require that $p_1 < \ldots < p_m$ in the order on $D \setminus \{p_0\}$ induced by the orientation. By $\mathcal{M}_{a;\mathbf{b};A}(V)$ we denote the moduli space of disks $u\colon \dot{D} \to \mathbb{R} \times Y$ which are J-holomorphic in the sense that

$$du + J du \circ i = 0$$

for some conformal structure $(\dot{D}, i)$, and such that u satisfies the following conditions.

- The boundary of u is on V and becomes a cycle in homology class $A \in H_1(V)$ when closed up with appropriately oriented capping paths.
- At the puncture p_0, u is is asymptotic to the Reeb chord a at $+\infty$.
- At the puncture p_i, where $i > 0$, u is asymptotic to the Reeb chord b_i at $-\infty$.

In general, we call punctures at which the disk is asymptotic to $+\infty$ *positive*, and punctures at which the disk is asymptotic to $-\infty$ *negative*.

Observe that, since $\mathbb{R} \times Y$ satisfies a convexity condition, there are no non-constant pseudo-holomorphic disks with boundary on V contained in a compact set.

Suppose that V is cylindrical outside of the set $[N_-, N_+] \times Y$. For any smooth disk D with the asymptotical behavior as above, we define its $d\lambda$-

and λ-*energy*, respectively, as follows

$$E_{d\lambda}(D) := \int_D d\lambda,$$

$$E_\lambda(D) := \sup_{\substack{\rho\colon (-\infty,N_-]\to\mathbb{R}_{\geq 0}\\ \int_{(-\infty,N_-]}\rho(t)dt=1}} \int_D \rho(t)dt\wedge\lambda + \sup_{\substack{\rho\colon [N_+,+\infty)\to\mathbb{R}_{\geq 0}\\ \int_{[N_+,+\infty)}\rho(t)dt=1}} \int_D \rho(t)dt\wedge\lambda.$$

The λ-energy of a pseudo-holomorphic disk is always positive. The $d\lambda$-energy of a pseudo-holomorphic disk is always non-negative, and is positive unless the disk is $\mathbb{R}$-invariant, that is, contained in a strip $\mathbb{R}\times a$ over some Reeb chord a. In the case when $V=\mathbb{R}\times L$, we have the formula

$$E_{d\lambda} = \ell(a) - \ell(b_1) - \ldots - \ell(b_m).$$

We define the *total energy* of a disk D by

$$E(D) := E_{d\lambda}(D) + E_\lambda(D) + \int_{D\cap([N_-,N_+]\times Y)} dt\wedge\lambda.$$

There is a version of Gromov's compactness theorem (see [Gro]) which applies to the setting of Symplectic Field Theory. It was introduced by Hofer and is valid for pseudo-holomorphic curves in symplectic manifolds with non-compact cylindrical ends. In particular, the results are valid in symplectizations with cylindrical almost complex structures.

The results in [BEH$^+$] imply that a sequence of pseudo-holomorphic disks in moduli spaces as above, satisfying a bound on their total energy, has a subsequence converging (in a certain sense) to a limit which is a *holomorphic building*. We refer to [BEH$^+$] for the exact statements and definitions. However, in our situation, a holomorphic building consists of a finite collection of holomorphic curves in $\mathbb{R}\times Y$, where each curve has an associated level $i\in\mathbb{Z}$, such that

- The curves at two consecutive levels have matching positive and negative asymptotics.
- The disks at level 0 have boundary on V.
- The disks at negative levels have boundary on L_1.
- The disks on positive levels have boundary on L_2.

One says that a layer is trivial if it consists of only $\mathbb{R}$-invariant solutions. A holomorphic building is allowed to consist of at most one trivial layer.

One says that an almost complex structure is *regular* if the above moduli spaces are transversely cut out. In this case, they are smooth manifolds satisfying

$$\dim \mathcal{M}_{a;\mathbf{b};A}(V) = |a| - |b_1| - \ldots - |b_m| + \mu(A).$$

Observe that, in the case when the almost complex structure is cylindrical, $\mathcal{M}_{a;\mathbf{b};A}(\mathbb{R}\times L)$ has a natural $\mathbb{R}$-action induced by the translation of the symplectization coordinate. Hence, in the case when the almost complex structure is regular, disks in such a 0-dimensional moduli spaces must be $\mathbb{R}$-invariant, and hence consist of trivial strips $\mathbb{R} \times a$ over Reeb chords.

Finally, it can be shown that the dimension formula is additive, that is, that the dimension of the disk converging to a holomorphic building is equal to the sum of the dimensions of the disks in the building.

2.3. **The differential.** For a Reeb-chord generator $a \in \mathcal{Q}(L)$, we define

$$\partial(a) := \sum_{\dim \mathcal{M}_{a;\mathbf{b};A}(\mathbb{R}\times L)=1} |\mathcal{M}_{a;\mathbf{b};A}(\mathbb{R} \times L)/\mathbb{R}| b_1 \cdot \ldots \cdot b_m,$$

where we assume that we have chosen a cylindrical almost complex structure J on $\mathbb{R} \times Y$ for which all moduli spaces in the definition are transversely cut out, and hence of the expected dimensions.

That this count is well-defined follows from the compactness theorem for the moduli spaces, since the disks in the above moduli spaces are of bounded total energy for any fixed a, together with the fact that there only are finitely many terms in the sum, which follows from the non-negativity of the $d\lambda$-action, since there only are finitely many Reeb chords below any given action. We extend the differential to all of $\mathcal{A}(L)$ via the Leibniz rule

$$\partial(ab) = \partial(a)b + a\partial(b).$$

The dimension-formula above shows that ∂ has degree -1.

The following standard argument shows that $\partial^2 = 0$. Consider the boundary $\partial\overline{\mathcal{M}}_{a;\mathbf{b};A}(\mathbb{R} \times L)/\mathbb{R}$ of the compactification of a 1-dimensional moduli space $\mathcal{M}_{a;\mathbf{b};A}(\mathbb{R} \times L)/\mathbb{R}$. By the additivity of dimensions, the boundary consist of buildings of two levels, where each level contains exactly one non-trivial solution coming from a moduli space occurring in the definition of ∂, and hence each boundary component contributes to the coefficient in front of $\mathbf{b}$ in the expression $\partial^2(a)$. Conversely, a gluing argument shows that each coefficient of a word contributing to $\partial^2(a)$ corresponds to a count of boundary points of such a 1-dimensional moduli space. Consequently, these coefficients are even, and thus vanishing.

Remark 2.1. The condition on λ that all periodic Reeb orbits are non-degenerate and have Conley-Zehnder index at least two is needed for the following reason. It implies that every pseudo-holomorphic plane asymptotic to a Reeb orbit in $\mathbb{R} \times Y$ is of formal dimension 2. When the moduli spaces of pseudo-holomorphic punctured disks are transversely cut-out, 1-dimensional families of such disks cannot split into a building containing a plane, since such a building would have to contain a disk of negative dimension by the additivity of the (formal) dimensions.

One defines the *Legendrian contact homology of* L to be the homology

$$HC_\bullet(L) := H_\bullet(\mathcal{A}(L), \partial).$$

2.4. **The invariance.**

The following theorem implies that the Legendrian contact homology is invariant under Legendrian isotopy.

Theorem 2.2 ([EES3], [Ekh2]). *Let $L \subset Y$ be a Legendrian submanifold. There is a regular cylindrical almost complex structure on $\mathbb{R} \times Y$ making the appropriate moduli spaces transversely cut out. The homotopy type of its associated DGA*

$$(\mathcal{A}(L), \partial)$$

is preserved under Legendrian isotopy and is independent of the choice of a generic cylindrical almost complex structure. In particular, $HC_\bullet(L)$ is invariant under Legendrian isotopy.

Remark 2.3. This statement follows from the more general Theorem 1.2 in [Ekh2]. Also, see Remark 3.9 in in [Ekh2].

The invariance proof in [Ekh2] is proved as follows. Given a Legendrian isotopy from L to L', one can construct an exact Lagrangian cobordism V_1 from L to L', and one exact Lagrangian cobordism V_2 from L' to L, such that the concatenation of V_1 with V_2 is Hamiltonian isotopic to the trivial cylinder over L, and the concatenation of V_2 with V_1 is Hamiltonian isotopic to the trivial cylinder over L'. The invariance then follows from the results below concerning exact Lagrangian cobordisms.

2.5. **Exact Lagrangian cobordisms and chain maps.**

Let V be an exact Lagrangian cobordism from L_1 to L_2. A standard construction in Symplectic Field Theory gives an induced DGA-morphism

$$\Phi\colon (\mathcal{A}(L_2), \partial_2) \to (\mathcal{A}(L_1), \partial_1),$$

which we now describe.

Let $\mathcal{M}_{a;\mathbf{b};A}(V)$ be the moduli space of punctured pseudo-holomorphic disks in $\mathbb{R} \times Y$ having boundary on V in homology class $A \in H_1(V)$, a positive puncture asymptotic to the Reeb chord a on L_2, negative punctures asymptotic to the Reeb chords $b_1, \dots, b_m$ on L_1, where $\mathbf{b} = b_1 \cdot \dots \cdot b_m$. The map Φ is defined on the Reeb chord-generator a of $\mathcal{A}(L_2)$ by the formula

$$\Phi(a) = \sum_{\dim \mathcal{M}_{a;\mathbf{b};A}(V)=0} |\mathcal{M}_{a;\mathbf{b};A}(V)| b_1 \cdot \dots \cdot b_m,$$

and then extended to an algebra map.

Again, the total energy is bounded for the curves in $\mathcal{M}_{a;\mathbf{b};A}(V)$, and the expression is well-defined by the above compactness theorem.

The fact that Φ is a chain-map follows by considering the boundary $\partial\overline{\mathcal{M}}_{a;\mathbf{b};A}(V)$ of the compactification of a 1-dimensional moduli space. It consist of holomorphic buildings of two levels, one which contains rigid solutions with boundary on V, and one which contains exactly one non-trivial 1-dimensional solution with boundary on either $\mathbb{R} \times L_1$ or $\mathbb{R} \times L_2$, and hence each boundary component contributes to the coefficient in front of $\mathbf{b}$ in the expression $\partial_1 \circ \Phi(a) + \Phi \circ \partial_2(a)$. Conversely, a gluing argument shows that each coefficient of a word in the expression $\partial_1 \circ \Phi(a) + \Phi \circ \partial_2(a)$ corresponds to a count of boundary points of such 1-dimensional moduli spaces. Hence, this expression vanishes and the chain-map property follows.

Theorem 2.4 ([Ekh2]). *Let $V \subset Y$ be an exact Lagrangian cobordism from the Legendrian submanifold L_1 to L_2. There is a regular cylindrical almost complex structure on $\mathbb{R} \times Y$ making the the appropriate moduli spaces transversely cut-out. The induced map*

$$\Phi\colon\ (\mathcal{A}(L_2), \partial_2) \to (\mathcal{A}(L_1), \partial_1)$$

is a DGA morphism whose homotopy-class is independent of the choice of a regular cylindrical almost complex structure and the (compactly supported) Hamiltonian isotopy class of V.

Remark 2.5. This follow from the more general statement [Ekh2, Lemma 3.15] together with the arguments in [Ekh2, Section 4].

The invariance under Hamiltonian isotopy of V and choice of cylindrical almost complex structure contains one substantial difficulty. Here one has to take into account pseudo-holomorphic disks of formal dimension -1, which arise in generic 1-parameter families of moduli spaces. We refer to [Ekh2, Section 4] for a treatment of this.

2.6. Augmentations.

An *augmentation* is a unital DGA-map

$$\varphi\colon\ (\mathcal{A}(L), \partial) \to \mathbb{Z}_2,$$

where $\mathbb{Z}_2$ is considered as a unital DGA with trivial differential. Augmentations obviously pull back by chain maps. It follows that if L_1 is exact Lagrangian cobordant to L_2, the existence of an augmentation for $(\mathcal{A}(L_1), \partial_1)$ implies the existence of an augmentation for $(\mathcal{A}(L_2), \partial_2)$.

The existence of an augmentation implies that one can define a linearized version of Legendrian contact homology. We refer to [Che] for a treatment of this.

2.7. Tame isomorphisms.

Let $\mathcal{A} = \mathbb{Z}_2\langle a_i \rangle_{i \in I}$ and $\mathcal{A}' = \mathbb{Z}_2\langle a_i' \rangle_{i \in I}$ be free unital algebras over $\mathbb{Z}_2$. An isomorphism

$$\varphi\colon\ \mathcal{A} \to \mathcal{A}'$$

of DGAs is *tame* if, after some identification of the generators of $\mathcal{A}$ and $\mathcal{A}'$, it can be written as a composition of *elementary* automorphisms, i.e. automorphisms defined on the generators of $\mathcal{A}$ by

$$\varphi(a_i) = \begin{cases} a_i & i \neq j \\ a_j + w & i = j \end{cases}$$

for some fixed j, where w is an element of the unital subalgebra generated by $\{a_i;\ \ i \neq j\}$.

3. The definition of a Legendrian ambient surgery

We define the notion of *Legendrian ambient surgery data* for a Legendrian submanifold $L \subset Y$. This data determines a Legendrian isotopy class of a Legendrian submanifold $L_S \subset Y$, where L_S is obtained by surgery on L along the framed sphere $S \subset L$. The construction will be called a *Legendrian ambient surgery* on L. Moreover, the construction provides an exact Lagrangian handle-attachment cobordism from L to L_S.

3.1. **Legendrian ambient surgery data.**

Let (Y, λ) be a contact manifold of dimension $2n+1$ with contact distribution $\xi := \ker \lambda$ and $L \subset Y$ a Legendrian submanifold, which hence satisfies $\dim L = n$. In the following, for a linear subspace $A \subset \xi$, we let $A^{d\lambda} \subset \xi$ denote the symplectic complement of A with respect to $d\lambda$, i.e. the subspace consisting of vectors $v \in \xi$ satisfying $d\lambda(A, v) = 0$. The following data, referred to as *Legendrian ambient surgery data*, is needed to specify a Legendrian embedding into Y of the manifold obtained by surgery on L.

(a) *The framed attaching sphere S.* An embedded k-sphere, where $k < n$, together with a choice of framing of its normal bundle $N \subset TL|_S$.

(b) *The bounding disk D_S of S.* An embedded isotropic $(k+1)$-disk $D_S \subset Y$ with boundary $\partial D_S = S$, interior disjoint from L, and an outward normal field G nowhere tangent to L which satisfies the following conditions. Let H be the vector-field in N defined uniquely up to homotopy by the requirement that $d\lambda(G, H) > 0$. We require that H is homotopic to a constant vector-field relative the above framing.

(c) *A compatible Lagrangian framing of the symplectic normal bundle N' of D_S.* From condition (b) it follows that the hyperplane field

$$T(D_S)^{d\lambda}|_S \cap N \subset N$$

along S has a natural framing induced by N, and defined uniquely up to homotopy. Let $N' \to D_S$ denote the symplectic normal bundle of D_S. Observe that $\mathbb{C} \otimes (T(D_S)^{d\lambda}|_S \cap N)$ is isomorphic to $N'|_S$ and hence we get an induced Lagrangian framing on $N'|_S$. We require that this Lagrangian framing extends to all of N', and we choose one such extension.

Remark 3.1. In the case when $k = 0$ condition (b) reduces to the requirement that the outward normal of D_S is not tangent to L. Moreover, any framing of $N'|_S$ in (c) extends to all of N' and, moreover, the choice of this extension lives in $\pi_1 U(n-1) \simeq \mathbb{Z}$.

3.2. **Topological conditions for the existence of Legendrian ambient surgery data in the subcritical case.**

The existence of a bounding disk and a compatible trivialization of the symplectic normal bundle, given a framed attaching sphere, can be described in the following topological terms in the case when $k < n - 1$, that is, in the subcritical case.

Suppose that we have a smoothly embedded disk $F\colon D^{k+1} \to Y$ with boundary coinciding with S and interior disjoint from L, such that the disk has an outward normal field G tangent to ξ and satisfying condition (b) above. To that end, we require

$$TS + \mathbb{R}G \subset \xi$$

to be an isotropic plane-field. We will still denote the disk by D_S, even though it is not necessarily isotropic in the interior.

Consider the Lagrangian framing of

$$\mathbb{C} \otimes (TS \oplus \mathbb{R}G \oplus (T(D_S)^{d\lambda}|_S \cap N)) \simeq \xi|_S$$

along S, where the summand $TS \oplus \mathbb{R}G$ is given the framing induced by $T(D_S)$, and where the last summand has the framing induced by N. We suppose that this Lagrangian framing of $\xi|_S$ can be extended to all of $\xi|_{D_S}$, and we choose one such extension.

The contact distribution ξ has a trivialization along D_S, and hence the obstruction to extending the above framing lives in $\pi_k U(n)$, while the choice of the homotopy-class of the extension lives in $\pi_{k+1} U(n)$.

Since $\dim D_S = k+1 \leq n-1$ by assumption, the h-principle for subcritical isotropic embeddings implies that D_S is isotopic relative its boundary to an embedded isotropic disk with tangent space homotopic to the sub-bundle of $\xi|_{D_S}$ spanned by the the first $k+1$ sections of the above framing. See [EM] for more details. The latter $n-k-1$ sections of the above framing induces a Lagrangian framing of the symplectic normal bundle of the bounding disk required by (c).

Remark 3.2. The above h-principle also implies that, in the case $k < n-1$, for topologically equivalent ambient surgery data there is an isotopy in $Y \setminus L$ between the corresponding isotropic bounding disks, which moreover is an isotopy through isotropic disks, and which induces the given homotopy of the Legendrian ambient surgery data. However, there is no such h-principle when $k = n-1$ since D_S is Legendrian in that case.

3.3. **The standard model for an exact Lagrangian handle attachment.**

Recall that the jet space $(J^1(M), dz+\theta)$ has the natural projections

$$\Pi_F\colon\ J^1(M) \to M \times \mathbb{R},$$
$$\Pi_L\colon\ J^1(M) \to T^*M,$$

called the *front projection* and the *Lagrangian projection*, respectively. A Legendrian submanifold of $J^1(M)$ is determined (up to a translation in the the Reeb-direction) by its image under either one of these projections.

The image of a Legendrian submanifold $L \subset J^1(M)$ under Π_L is an exact immersed Lagrangian submanifold of the exact symplectic manifold $(T^*M, d\theta)$. Reeb chords on L correspond to self-intersections of Π_L. In the front projection, Reeb chords correspond to two sheets of $\Pi_F(L)$ having a common tangency above a given point in the base M.

We will use the above projections to construct a Legendrian standard model $L_{0,k} \subset J^1(\mathbb{R}^n)$ of a neighborhood of a framed k-sphere $S_{0,k} \subset L_{0,k}$, and a Legendrian submanifold $L_{\epsilon,k} \subset J^1(\mathbb{R}^n)$ which is obtained from $L_{0,k}$ by surgery along $S_{0,k}$. We also construct an exact Lagrangian cobordism $W_\epsilon \subset \mathbb{R} \times J^1(\mathbb{R}^n)$ from $L_{0,k}$ to $L_{\epsilon,k}$ which is diffeomorphic to the manifold obtained by a $(k+1)$-handle attachment on $(-\infty,-1] \times L_{0,k}$ along

$$S_{0,k} \subset L_{0,k} = \partial((-\infty,-1] \times L_{0,k}).$$

Let t be the coordinate of the $\mathbb{R}$-factor in the symplectization $\mathbb{R} \times J^1(\mathbb{R}^n)$, and let $(\mathbf{q},\mathbf{p},z)$ be canonical coordinates on $\left(J^1(\mathbb{R}^n), dz - \sum_{i=1}^n p_i dq_i\right)$. Making the coordinate change

$$\begin{cases} t = \log x_{n+1}, \\ z = y_{n+1}, \\ p_i = y_i / x_{n+1}, \\ q_i = x_i, \end{cases}$$

we identify the symplectization $\left(\mathbb{R} \times J^1(\mathbb{R}^n), d\left(e^t\left(dz - \sum_{i=1}^n p_i dq_i\right)\right)\right)$ with the exact symplectic manifold

$$(T^*(\mathbb{R}^n \times \mathbb{R}), d(-y_1 dx_1 - \ldots - y_n dx_n + x_{n+1} dy_{n+1}))$$

endowed with the coordinates

$$(\mathbf{x},\mathbf{y}) = (x_1, \ldots, x_{n+1}, y_1, \ldots, y_{n+1}),$$

where $\mathbf{x}$ form standard coordinates on the base $\mathbb{R}^n$ and $\mathbf{y}$ form coordinates induced by the coframe $\{dx_i\}$. The above identification is an exact symplectomorphism.

An exact Lagrangian submanifold of $\mathbb{R} \times J^1(\mathbb{R}^n)$ can thus be identified with an exact Lagrangian submanifold of $T^*(\mathbb{R}^n \times \mathbb{R})$ with respect to the above primitive of the symplectic form. Observe that such an exact Lagrangian submanifold is also exact with respect to the Liouville form on $T^*(\mathbb{R}^n \times \mathbb{R})$ and, hence, it admits a Legendrian lift to $J^1(\mathbb{R}^n \times \mathbb{R})$.

In conclusion, an exact Lagrangian submanifold of $\mathbb{R} \times J^1(\mathbb{R}^n)$ can be represented by a front over the base of $T^*(\mathbb{R}^n \times \mathbb{R})$ via the identification

$$y_i = \partial_{x_i} f(\mathbf{x}),$$

using the above coordinates, where the graph of $f(\mathbf{x})$ determines a sheet of the front projection of its Legendrian lift. A primitive of the pull-back of the form

$$-y_1 dx_1 - \ldots - y_n dx_n + x_{n+1} dy_{n+1}$$

to the exact Lagrangian submanifold is given by

$$-f(\mathbf{x}) + x_{n+1} y_{n+1} + C.$$

Under the above identification, the property of being cylindrical over a Legendrian submanifold in $J^1(\mathbb{R}^n)$ for a Lagrangian submanifold of $T^*\mathbb{R}^{n+1} \simeq$

$\mathbb{R} \times J^1(\mathbb{R}^n)$, is equivalent to the sheets of the front projection of its Legendrian lift being given by functions on the form

$$f(\mathbf{x}) = g(x_1, \ldots, x_n)x_{n+1}$$

above a region in the base of $T^*(\mathbb{R}^n \times \mathbb{R})$.

3.3.1. *The model for an exact Lagrangian handle attachment W_ϵ.* We now define the cobordism W_ϵ. Fix a real number $\eta > 0$. For $\epsilon > 0$, let

$$\rho_\epsilon \colon \mathbb{R}_{\geq 0} \to [0,1]$$

be a smooth non-increasing function satisfying the following.

- $\rho_\epsilon(0) = 1$.
- $\rho_\epsilon(x^2) = 0$ when $x^2 \geq \eta + (2/3)\epsilon$.
- $\frac{d^2}{dx^2}\left(x^2 + \rho_\epsilon\left(x^2\right)(\eta + \epsilon/2)\right) > 0$ for all $x \in \mathbb{R}$.

Furthermore, let

$$\sigma_\epsilon \colon \mathbb{R} \to \mathbb{R}$$

be a smooth non-decreasing function satisfying the following.

- $0 \leq \sigma_\epsilon'(x) \leq (\eta + \epsilon)\epsilon^{-1/3}$.
- $\sigma_\epsilon(x) = 0$ for $x \leq 1 - \epsilon^{1/3}$.
- $\sigma_\epsilon(x) = \eta + \epsilon/2$ for $x \geq 1 + \epsilon^{1/3}$.
- $\sigma_\epsilon(1) = \eta$, $\sigma_\epsilon'(1) > 0$.

Finally, we define

$$\varphi_\epsilon \colon \mathbb{R}^{n+1} \to \mathbb{R}$$

by

$$\varphi_\epsilon(\mathbf{x}) = \rho_\epsilon\left(x_1^2 + \ldots + x_{k+1}^2\right)\sigma_\epsilon\left(x_{n+1}\right).$$

We now consider the front over $\mathbb{R}^{n+1}$ corresponding to the graphs of the two functions $\pm F_\epsilon(\mathbf{x})x_{n+1}$, where

$$F_\epsilon(\mathbf{x}) := \left(\left(x_1^2 + \ldots + x_{k+1}^2\right) - \left(x_{k+2}^2 + \ldots + x_n^2\right) + \varphi_\epsilon(\mathbf{x}) - \eta\right)^{3/2},$$

above the set

$$\left\{ \begin{array}{l} \left(x_1^2 + \ldots + x_{k+1}^2\right) - \left(x_{k+2}^2 + \ldots + x_n^2\right) + \varphi_\epsilon(\mathbf{x}) \geq \eta, \\ x_{n+1} > 0 \end{array} \right\},$$

and denote the corresponding Lagrangian submanifold in $T^*\mathbb{R}^{n+1}$ by W_ϵ.

Observe that W_ϵ can be identified with an exact Lagrangian submanifold of $\mathbb{R} \times J^1(\mathbb{R}^n)$ which is cylindrical over a Legendrian submanifold above the complement of

$$\left\{ \begin{array}{l} x_1^2 + \ldots + x_{k+1}^2 \leq \eta + \epsilon, \\ 1 - \epsilon^{1/3} \leq x_{n+1} \leq 1 + \epsilon^{1/3} \end{array} \right\} \subset \mathbb{R}^n \times \mathbb{R},$$

and that the primitive of the pull-back of the form

$$-y_1dx_1 - \ldots - y_ndx_n + x_{n+1}dy_{n+1}$$

may be taken to be zero there.

We also consider the exact Lagrangian submanifold W_0 defined by the graphs of

$$\pm\left((x_1^2+\ldots+x_{k+1}^2)-(x_{k+2}^2+\ldots+x_n^2)-\eta\right)^{3/2}x_{n+1}$$

above the set

$$\left\{\begin{array}{l}(x_1^2+\ldots+x_{k+1}^2)-(x_{k+2}^2+\ldots+x_n^2)\geq\eta,\\ x_{n+1}>0\end{array}\right\}.$$

Obviously, W_0 is cylindrical over a Legendrian submanifold, and the primitive of the pull-back of the form

$$-y_1dx_1-\ldots-y_ndx_n+x_{n+1}dy_{n+1}$$

may be taken to vanish.

By

$$L_{0,k}\subset J^1(\mathbb{R}^n)$$

and

$$L_{\epsilon,k}\subset J^1(\mathbb{R}^n)$$

we denote the Legendrian submanifolds over which W_ϵ is cylindrical at its negative and positive end, respectively. Clearly, W_0 is a cylinder over $L_{0,k}$.

Observe that $L_{\epsilon,k}$ is diffeomorphic to the manifold obtained by performing surgery on $L_{0,k}$ along the framed k-sphere

$$S_{0,k}:=\left\{\begin{array}{l}q_1^2+\ldots+q_{k+1}^2=\eta,\\ q_{k+2}=\ldots=q_n=p_1=\ldots=p_n=z=0\end{array}\right\}$$

with framing

$$N_{0,k}:=\langle q_1\partial_{p_1}+\ldots+q_{k+1}\partial_{p_{k+1}},\partial_{q_{k+2}},\ldots,\partial_{q_n}\rangle.$$

The cobordism W_ϵ is diffeomorphic to the manifold obtained by a handle-attachment on $(-\infty,-1]\times L_{0,k}$ along

$$S_{0,k}\subset L_{0,k}=\partial((-\infty,-1]\times L_{0,k}).$$

By applying Formula 2.2, one sees that the Maslov class vanishes for W_ϵ, and hence for both $L_{0,k}$ and $L_{\epsilon,k}$.

3.3.2. *The model for the Legendrian ambient surgery data.* The Legendrian submanifold $L_{0,k}$ supports the Legendrian ambient surgery data consisting of the framed sphere $S_{0,k}$ defined above, the isotropic bounding disk

$$D_{0,k}:=\left\{\begin{array}{l}q_1^2+\ldots+q_{k+1}^2\leq\eta,\\ q_{k+2}=\ldots=q_n=p_1=\ldots=p_n=z=0\end{array}\right\},$$

and the compatible Lagrangian framing

$$\{\partial_{q_{k+2}},\ldots,\partial_{q_n}\}$$

of the symplectic normal bundle

$$N'_{0,k}:=\mathbb{C}\langle\partial_{q_{k+2}},\ldots,\partial_{q_n}\rangle$$

of $D_{0,k}$.

3.3.3. *The Reeb chords on $L_{0,k}$ and $L_{\epsilon,k}$.* Note that $L_{0,k}$ has no Reeb-chords, and that $L_{\epsilon,k}$ has exactly one Reeb chord c_{new} above $\mathbf{q}=0$. One computes

$$\ell(c_{new}) = \epsilon^{3/2}/\sqrt{2}.$$

Furthermore, this Reeb chord satisfies

$$|c_{new}| = n-k-1,$$

which can be seen by applying Formula 2.1. To that end, we observe that the difference-functions of the two sheets of the front has Morse index $n-(k+1)$ at the origin, and that the obvious capping path of c_{new} traverses exactly one cusp-edge in downward direction.

3.4. **The exact Lagrangian handle attachment induced by the Legendrian ambient surgery data.** Suppose that we are given Legendrian ambient surgery data for the Legendrian submanifold $L \subset Y$ consisting of a k-sphere $S \subset L$ with framed normal bundle N and an isotropic bounding $(k+1)$-disk $D_S \subset Y$ with a compatible Lagrangian framing of its symplectic normal bundle N'.

We now construct an exact Lagrangian cobordism $V_\epsilon \subset \mathbb{R} \times Y$ from L to L_S such that V_ϵ is diffeomorphic to a $(k+1)$-handle attachment and L_S is obtained from L by k-surgery along S with the above framing on N.

3.4.1. *A standard neighborhood of the isotropic bounding disk.* Given the above data, we want to find coordinates in a neighborhood of D_S identifying it with a neighborhood of the disk $D_{0,k}$ defined in Section 3.3 such that L is identified with $L_{0,k}$ and, moreover, $L_{\epsilon,k}$ and $L_{0,k}$ coincide outside an even smaller neighborhood, for $\epsilon > 0$ small enough.

Let H be a choice (which is unique up to homotopy) of a vector-field in N which satisfies $d\lambda(G,H) > 0$ for the outward normal vector-field G of D_S along the boundary.

The normal form of isotropic subspaces of contact manifolds (see [McD]) implies that there is a contactomorphism which identifies a neighborhood of $D_S \subset Y$ with some neighborhood of the isotropic $(k+1)$-disk

$$D_{0,k} \subset J^1(\mathbb{R}^n)$$

such that the Lagrangian framing of the symplectic normal bundle N' of D_S is identified with the Lagrangian framing

$$\{\partial_{q_{k+2}}, \dots, \partial_{q_n}\}$$

of the symplectic normal bundle $N'_{0,k}$ of $D_{0,k}$. Moreover, the contactomorphism can be taken to preserve the contact form.

Since

$$TL|_S \simeq TS \oplus \mathbb{R}H \oplus \mathbb{R}\langle \partial_{q_{k+2}}, \dots, \partial_{q_n} \rangle,$$

we may, after possibly choosing smaller neighborhoods and perturbing L with a small Legendrian isotopy, assume that the contactomorphism identifying D_S with $D_{0,k}$, and hence S with $S_{0,k}$, moreover satisfies the following:

- The framing of the normal bundle N of S coincides with the above framing of $N_{0,k}$ defined in Section 3.3.
- A neighborhood of $S \subset L$ is identified with a neighborhood of $S_{0,k} \subset L_{0,k}$.

3.4.2. *The constructions of V_ϵ and L_S.* The identification of a neighborhood of D_S with a neighborhood of $D_{0,k}$ via the contact-form preserving map constructed above induces an exact symplectomorphism of a neighborhood of

$$\mathbb{R} \times D_S \subset \mathbb{R} \times Y$$

with a neighborhood of

$$\mathbb{R} \times D_{0,k} \subset \mathbb{R} \times J^1(\mathbb{R}^n)$$

which, moreover, preserves the symplectization coordinate t, and identifies $\mathbb{R} \times L$ with W_0.

We make the identification of $\mathbb{R} \times J^1(\mathbb{R}^n)$ with $T^*\mathbb{R}^{n+1}$ as in Section 3.3. After choosing a small enough $\epsilon > 0$, we may thus assume that the neighborhood

$$U_\epsilon := \left\{ \begin{array}{l} x_1^2 + \ldots + x_{k+1}^2 < \eta + \epsilon, \\ x_{k+2}^2 + \ldots + x_n^2 < 2\epsilon, \\ x_{n+1} > 0 \\ |y_i| < 3\sqrt{\epsilon(\eta+\epsilon)}x_{n+1}, \ \ i \leq n, \\ |y_{n+1}| < 3\epsilon^{1/6}(\eta + \epsilon) \end{array} \right\} \subset T^*\mathbb{R}^{n+1}.$$

of $\mathbb{R} \times D_{0,k}$ is contained in the image of the above contactomorphism.

Lemma 3.3. *For $\epsilon > 0$ small enough, W_ϵ and W_0 coincide outside of the open set $U_{\epsilon'}$ for any $\epsilon' \geq (2/3)\epsilon$.*

Proof. We show the claim for $U_{(2/3)\epsilon}$, that is, we set $\epsilon' = (2/3)\epsilon$. The general statement obviously follows from this.

Consider the function

$$f_y(x) := x^2 + \rho_\epsilon(x^2)y,$$

where ρ is as defined in Section 3.3. Recall that $f_y(x)$ is symmetric in x and that $f_y(x) = x^2$ whenever $x^2 \geq \eta + \epsilon'$. Since $f_0'' > 0$ obviously holds, and $f_{\eta+\epsilon/2}'' > 0$ holds by construction, we conclude that $f_y'' > 0$ for all $0 \leq y \leq \eta + \epsilon/2$. From this we conclude that

$$f_y(0) \leq f_y(x) < \eta + \epsilon'$$

for all $x^2 < \eta + \epsilon'$, $0 \leq y \leq \eta + \epsilon/2$. Hence we get the bound

$$0 \leq f_y(0) \leq x^2 + \rho_\epsilon\left(x^2\right)y - \eta < \epsilon'$$

for all $x^2 < \eta + \epsilon'$ and $0 \leq y \leq \eta + \epsilon/2$.

The above inequality implies that the set

$$W_\epsilon \cap \left\{x_1^2 + \ldots + x_{k+1}^2 < \eta + \epsilon'\right\}$$

is contained inside

$$\left\{x_{k+2}^2+\ldots+x_n^2<\epsilon'\right\}$$

and, furthermore, that the inequality

$$F_\epsilon(\mathbf{x})<(\epsilon')^{3/2}$$

holds on the same set.

Furthermore, since $\rho_\epsilon(x^2)$ vanishes when $x^2\geq\eta+\epsilon'$, it follows that W_ϵ and W_0 coincide outside of the set

$$\left\{x_1^2+\ldots+x_{k+1}^2<\eta+\epsilon'\right\},$$

and by the above remark, outside of

$$O:=\left\{\begin{array}{l} x_1^2+\ldots+x_{k+1}^2<\eta+\epsilon', \\ x_{k+2}^2+\ldots+x_n^2<\epsilon' \end{array}\right\}.$$

Since $f_y(x)$ is symmetric in x and has increasing derivative by assumption, and since $f'_y(\pm\sqrt{\eta+\epsilon'})=\pm2\sqrt{\eta+\epsilon'}$, we have the bounds

$$\left|\frac{\partial}{\partial x_i}F_\epsilon(\mathbf{x})\right|\leq(3/2)|F_\epsilon(\mathbf{x})^{1-2/3}|\left(2\sqrt{\eta+\epsilon'}\right)<3\sqrt{\epsilon'}\sqrt{(\eta+\epsilon')}$$

for $1\leq i\leq k+1$, and

$$\left|\frac{\partial}{\partial x_i}F_\epsilon(\mathbf{x})\right|\leq(3/2)|F_\epsilon(\mathbf{x})^{1-2/3}|\left(2\sqrt{\epsilon'}\right)<3\sqrt{\epsilon'}\sqrt{(\eta+\epsilon')}$$

for $k+2\leq i\leq n$. inside O, which implies that $W_\epsilon\cap O$ is contained in the set

$$\left\{|y_i|<3\sqrt{\epsilon'(\eta+\epsilon')}x_{n+1},\ \ 1\leq i\leq n\right\}.$$

Finally, since $0\leq\sigma'_\epsilon(x)\leq(\eta+\epsilon)\epsilon^{-1/3}$, together with the estimate $F_\epsilon(\mathbf{x})<(\epsilon')^{3/2}$ on O, we get the bound

$$\begin{aligned}&\left|\frac{\partial}{\partial x_{n+1}}F_\epsilon(\mathbf{x})x_{n+1}\right|\\ \leq\ \ & F_\epsilon(\mathbf{x})+\frac{3}{2}F_\epsilon(\mathbf{x})^{1-2/3}x_{n+1}\sigma'_\epsilon(x_{n+1})<(\epsilon')^{3/2}+\frac{3}{2}(\epsilon')^{1/2}\epsilon^{-1/3}(\eta+\epsilon).\end{aligned}$$

on O. For $\epsilon>0$ small enough, this implies that $W_\epsilon\cap O$ is contained inside

$$\left\{|y_{n+1}|<3(\epsilon')^{1/6}(\eta+\epsilon')\right\}.$$

In conclusion, W_ϵ and W_0 coincide outside of the set $U_{\epsilon'}$. □

Since W_0 and W_ϵ in particular coincide outside of the set U_ϵ, we can replace W_0 with W_ϵ inside U_ϵ which, after identifying U_ϵ with the neighborhood of $\mathbb{R}\times D_S$ as above, produces an exact Lagrangian cobordism V_ϵ inside $\mathbb{R}\times Y$ from L to a Legendrian submanifold diffeomorphic to the manifold obtained by k-surgery along S. We denote this Legendrian submanifold by $L_S\subset Y$.

Definition 3.4. Given Legendrian ambient surgery data for $L \subset Y$, the *Legendrian ambient surgery on* L is the Legendrian submanifold $L_S \subset Y$ defined as above, by replacing L with the standard model $L_{0,k}$ in a neighborhood of $D_S \subset Y$, where the identification is determined by the Legendrian ambient surgery data as above.

Remark 3.5. Observe that for $\epsilon > 0$ small enough the Legendrian isotopy class of $L_S \subset Y$ is independent of the homotopy class of the choices made for the Legendrian ambient surgery data and of the choices made in the above construction. Also, see Remark 3.2.

Remark 3.6. When $k < n-1$, the isotropic bounding disk D_S is subcritical and hence generically has no Reeb chords. In particular, for any $E > 0$ we may assume, after choosing $\epsilon > 0$ small enough, that the Reeb chords on L_S having action less than E consist of Reeb chords coinciding with the Reeb chords on L having action less than E, together with the Reeb chord c_{new} on L_S situated in the middle of the handle.

3.5. The effect of a Legendrian ambient surgery on the Maslov class.

The Maslov class of a Lagrangian cobordism pulls back to the Maslov classes of its Legendrian ends under the inclusion maps. Hence

$$\mu(H_1(L)), \mu(H_1(L_S)) \subset \mu(H_1(V_\epsilon)) \subset \mathbb{Z}.$$

Let $X \cup (D^{k+1} \times D^{n-k})$ denote a $(k+1)$-handle attachment on the manifold X, where X has non-empty boundary. From the associated long exact sequence in singular homology

$$\begin{aligned} \ldots \to H_2(D^{k+1} \times D^{n-k}, S^k \times D^{n-k}) \to H_1(X) \to \\ \to H_1(X \cup (D^{k+1} \times D^{n-k})) \to H_1(D^{k+1} \times D^{n-k}, S^k \times D^{n-k}) \to \\ \to H_0(X) \to \ldots, \end{aligned}$$

together with the fact that

$$H_1(D^{k+1} \times D^{n-k}, S^k \times D^{n-k}) \simeq H_1(S^{k+1})$$

we conclude that the above map

$$H_1(X) \to H_1(X \cup (D^{k+1} \times D^{n-k}))$$

is surjective except possibly in the case $k = 0$.

Setting $X = (-\infty, -1] \times L$ and identifying $X \cup (D^{k+1} \times D^{n-k})$ with V_ϵ, we conclude that

$$\mu(H_1(L)) = \mu(H_1(V_\epsilon)) \subset \mathbb{Z}$$

whenever L_S is obtained from L by an ambient k-surgery with $k > 0$.

Since, topologically, V_ϵ is obtained from a $(n-k)$-handle attachment on $(-\infty, -1] \times L_S$, the same argument shows that

$$\mu(H_1(L_S)) = \mu(H_1(L)) = \mu(H_1(V_\epsilon)) \subset \mathbb{Z}$$

whenever L_S is obtained from L by an ambient k-surgery with $0 < k < n-1$.

In the case $k = 0$, we made the observation that we have a choice living in

$$\pi_1(U(n-1)) \simeq \mathbb{Z}$$

for the Lagrangian framing of the symplectic normal bundle of the bounding 1-disk. In the case when the 1-handle attachment adds a generator in H_1, these choices correspond to the Maslov index of the newly added generator, which hence can be chosen arbitrarily.

4. The DGA of L twisted by $S \subset L$

Let $L \subset (Y, \lambda)$ be a chord-generic Legendrian submanifold of a contact manifold of dimension $2n + 1$, together with an embedded framed k-sphere $S \subset L$ which is disjoint from the Reeb chords on L. Here we make the requirement that $k < n - 1$.

Fix any number $E > 0$. In this section we will only consider the Reeb chords on L having action less than E. Thus, there are finitely many such Reeb chords and, moreover, they all are at some positive distance away from S.

In the case when L has only finitely many Reeb chords, we may of course take $E > 0$ big enough to capture all of them, however, for the general case, we refer to Section 6 for a direct limit construction which takes into account all of the Reeb chords.

4.1. The algebra.

We define the *Legendrian contact homology DGA of L twisted by S* as follows. As an algebra it is freely generated over $\mathbb{Z}_2$ by

$$\mathcal{A}(L, S)^E := \mathbb{Z}_2 \langle \mathcal{Q}(L)_E \cup \{s\} \rangle,$$

where $\mathcal{Q}(L)_E$ denotes the set of Reeb chords on L of action less than E, and where s is a formal variable. The Reeb-chord generators are graded as in Section 2.1, while we grade the formal variable by

$$|s| = n - k - 1.$$

In the case when $\mathcal{Q}(L)_E$ contains all the Reeb chords on L, we drop E from the notation and simply write $\mathcal{A}(L, S) := \mathcal{A}(L, S)^E$.

Remark 4.1. Note that in the case when there are Legendrian ambient surgery data for a surgery along S, the grading of s coincides with the grading of the new Reeb chord c_{new} which appears in the middle of the handle produced by the surgery (see Section 3.4).

4.2. The relevant moduli spaces.

We fix an identification of a neighborhood of $S \subset L$ with $S \times \mathbb{R}^{n-k}$ using the framing, and choose a non-zero vector $\mathbf{v} \in \mathbb{R}^{n-k} \setminus \{0\}$, which is considered as a non-vanishing normal vector field to $S \subset L$. We extend $\mathbf{v}$ to an $\mathbb{R}$-invariant vector field normal to $\mathbb{R} \times S \subset \mathbb{R} \times L$.

Let $a \in \mathcal{Q}(L)_E$ be a Reeb chord on L, $\mathbf{b} = b_1 \cdot \ldots \cdot b_m$ a word of Reeb chords on L, $\mathbf{w} = (w_1, \ldots, w_{m+1}) \in (\mathbb{Z}_{\geq 0})^{m+1}$, and $A \in H_1(L)$. In Section 8 we define the moduli spaces

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R} \times L, \mathbb{R} \times S, \mathbf{v})$$

which count pseudo-holomorphic disks $u \colon \dot{D} \to \mathbb{R} \times Y$, for any choice of conformal structure on $\dot{D} = D \setminus \{p_0, \ldots, p_m\}$, satisfying

- $u(\partial D^2) \subset \mathbb{R} \times L$ and, after closing up the boundary with appropriately oriented capping paths, one obtains $A \in H_1(L)$.
- u has a positive puncture asymptotic to a at p_0.
- u has negative punctures asymptotic to b_i at p_i for $i > 0$.
- u maps a number w_i of boundary points on the arc between p_i and p_{i+1} to $\mathbb{R} \times S$ (where we set $p_{m+1} := p_0$), such that each point is counted with multiplicity $l+1$, where l is the order of tangency of the boundary to $\exp_{\mathbb{R}\times S}(\mathbb{R}\mathbf{v})$ in case when the tangent of the boundary has a positive component in the $\mathbf{v}$-direction, and where otherwise $l = 0$.

For convenience, we will in the following denote these spaces by

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v}) := \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R} \times L, \mathbb{R} \times S, \mathbf{v}).$$

Their expected dimensions are given by the formula

$$\dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v}) = |a| - |b_1| - \ldots - |b_m| - |s|(w_1 + \ldots + w_{m+1}) + \mu(A),$$

as shown in Section 8.8. Observe that for cylindrical almost complex structures these spaces carry a natural $\mathbb{R}$-action induced by the translation in the symplectization direction.

We will also need the moduli spaces

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v}) := \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R} \times L, \mathbb{R} \times S, \mathbf{v}),$$

which are constructed in Section 9, where $\delta > 0$ and

$$\mathbf{k} = (k_1, \ldots, k_{w_1+\ldots+w_{m+1}}) \in (\mathbb{Z}_{\geq 0})^{w_1+\ldots+w_{m+1}}$$

is a tuple of increasing integers. These moduli spaces are defined as the above, but where the last condition is replaced by

- u maps a number w_i of boundary points on the arc between p_i and p_{i+1} (where we set $p_{m+1} := p_0$) to the parallel copies $\exp_{\mathbb{R}\times S}(k_i \delta \mathbf{v})$ of $\mathbb{R} \times S$, such that the i:th boundary point with respect to the order on $\partial D \setminus \{p_0\}$ induced by the orientation is mapped to $\exp_{\mathbb{R}\times S}(k_i \delta \mathbf{v})$. Here exp denotes the exponential map on $\mathbb{R} \times L$ for some choice of metric.

In the case when $\mathbf{k} = (0, 1, \ldots, w_1 + \ldots + w_{m+1} - 1)$, we will use the notation

$$\mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v}) := \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v}).$$

Lemma 4.2. *There is a cylindrical almost complex structure on* $\mathbb{R} \times Y$ *for which all moduli spaces*

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v})$$

are transversely cut out, as well as the moduli spaces

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v})$$

of expected dimension 1.

Proof. The goal is to apply Proposition 8.5, which requires that the complex structure is integrable in a neighborhood of $\mathbb{R} \times S$ and that S, L and $\mathbf{v}$ satisfy real-analyticity conditions in this neighborhood. Proposition 9.1 then ensures that we moreover can assume that the spaces

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v})$$

are transversely cut out for this almost complex structure, since the Reeb chords of interest are at some positive distance from $\mathbb{R} \times S$.

As in Section 3.4.1, we may use the theorem of the normal form for Legendrian submanifolds to show the following. Let the Legendrian submanifold $L_{0,k} \subset J^1(\mathbb{R}^n)$ and the sphere $S_{0,k} \subset L_{0,k}$ be as defined in Section 3.3. We can make an identification of a neighborhood U of $S \subset Y$ with a neighborhood of $S_{0,k} \subset J^1(\mathbb{R}^n)$ by a contact-form preserving contactomorphism for which:

- S is identified with $S_{0,k}$.
- $U \cap L$ is identified with a neighborhood of $S_{0,k} \subset L_{0,k}$.
- The vector field $\mathbf{v}$ along S is identified with the vector-field
$$q_1\partial_{p_1} + \ldots + q_{k+1}\partial_{p_{k+1}}$$
along $S_{0,k}$.

This contactomorphism induces an exact symplectomorphism identifying the neighborhood $\mathbb{R} \times U$ of

$$\mathbb{R} \times S \subset \mathbb{R} \times Y$$

with a neighborhood of

$$\mathbb{R} \times S_{0,k} \subset \mathbb{R} \times J^1(\mathbb{R}^n),$$

while preserving the symplectization coordinate.

We now choose a cylindrical almost complex structure on $\mathbb{R} \times Y$ which is integrable in the neighborhood $\mathbb{R} \times U$. Using the above identification of $\mathbb{R} \times U$ with a neighborhood in $\mathbb{R} \times J^1(\mathbb{R}^n)$, we there require the almost complex structure to be given by

$$\begin{aligned} J\partial_{q_i} &= \partial_{p_i} + y\partial_t, \\ J\partial_{p_i} &= -\partial_{q_i} - y\partial_z, \\ J\partial_t &= \partial_z, \\ J\partial_z &= -\partial_t. \end{aligned}$$

This cylindrical almost complex structure is integrable since its corresponding Nijenhuis tensor $N_J(X,Y)$ vanishes identically, as can be checked by a direct calculation, where on local sections X, Y of $T\mathbb{R}^{n+1}$ we have

$$N_J(X,Y) = [X,Y] + J([JX,Y] + [X,JY]) - J[X,Y].$$

We may identify the neighborhood $\mathbb{R} \times U$ with a neighborhood in $T^*\mathbb{R}^{n+1}$ as in Section 3.3. Under this identification, it immediately follows that the projections

$$(x_i, y_i/x_{n+1})\colon\ T^*\mathbb{R}^{n+1} \to \mathbb{C},\ \ i = 1, \dots, n$$

are holomorphic for the above cylindrical almost complex structure.

Finally, using the holomorphic projection

$$\mathbb{R} \times J^1(\mathbb{R}^n) \to T^*\mathbb{R}^n \simeq \mathbb{C}^n,$$

one can see that

$$\mathbb{R} \times S_{0,k} \subset \mathbb{R} \times L_{0,k} \subset \mathbb{R} \times J^1(\mathbb{R}^n)$$

both are real-analytic submanifolds, and that $\mathbf{v}$ is a real-analytic vector field. It follows that there is a holomorphic identification of a neighborhood of $\mathbb{R} \times S \subset \mathbb{R} \times Y$ with a neighborhood in

$$T^*(\mathbb{R} \times S^k \times \mathbb{R} \times \mathbb{R}^{n-k-1}) \subset T^*(\mathbb{R} \times \mathbb{R}^{k+1} \times \mathbb{R}^{n-k-1}),$$

where the latter spaces are endowed with the standard integrable complex structure, and where:

- $\mathbb{R} \times L$ is identified with the zero-section.
- $\mathbb{R} \times S$ is identified with the submanifold $\mathbb{R} \times S^k \times \{0\}$ of the zero-section.
- The vector-field $\mathbf{v}$ along $\mathbb{R} \times S$ is identified with the coordinate vector-field of the second $\mathbb{R}$-factor in the zero-section $\mathbb{R} \times S^k \times \mathbb{R} \times \mathbb{R}^{n-k-1}$.

It follows that Proposition 8.5 applies, which finishes the proof. □

4.3. **The boundary map.**

We define the differential of the Legendrian contact homology algebra $(\mathcal{A}(L,S)^E, \partial_{S,N})$ twisted by S as follows.

$$\partial_{S,N}(s) := 0,$$

$$\partial_{S,N}(a) := \sum_{\dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v})=1} |\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v})/\mathbb{R}| s^{w_1} b_1 s^{w_2} \cdot \ldots \cdot s^{w_m} b_m s^{w_{m+1}},$$

where $a \in \mathcal{Q}(L)_E$. We extend $\partial_{S,N}$ to $\mathcal{A}(L,S)^E$ using the Leibniz rule. From the above dimension formula it immediately follows that $\partial_{S,N}$ has degree -1.

According to the energy formula for pseudo-holomorphic disks with punctures, the above sum contains only finitely many terms. Furthermore, according to the compactness result in Proposition 8.6, there are only finitely many solutions in each rigid moduli space. Finally, by the compactness theorem in [BEH$^+$], the moduli space $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v})$, where a and $\mathbf{b}$ are fixed, is non-empty only for finitely many values of $A \in H_1(L)$. It follows that $\partial_{S,N}$ is well-defined.

Remark 4.3. The above complex is supposed to model the Legendrian contact homology DGA of a Legendrian submanifold L_S obtained by Legendrian ambient surgery on L along S, in the case when it can be performed. The formal generator s is supposed to correspond to the Reeb chord c_{new} produced by such a surgery. The following calculation can be seen as a motivation for why we should have $\partial_{S,N}(s) = 0$.

Since the Reeb chord c_{new} on L_S produced by a Legendrian ambient surgery can be made to have arbitrary small action, we immediately get that

$$\partial c_{new} = 0$$

in the case when $|c_{new}| \neq 1$, i.e. when $k \neq n - 2$.

In the case when $k = n - 2$ one can check that the same computation is true in the standard model $L_{\epsilon,k} \subset J^1(\mathbb{R}^n)$ by using the technique of gradient flow trees defined in [Ekh1]. In this case, there are exactly two rigid flow trees having a positive puncture at c_{new} and no negative punctures. This calculation is made precise in Lemma 5.6.

Remark 4.4. The differential will in general depend on the framing of S.

Proof of Theorem 1.2. We have already discussed the well-definedness of $\partial_{S,N}$. We start by showing ${\partial_{S,N}}^2 = 0$. Suppose that

$$\dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v}) = 2.$$

By the compactness results in [BEH$^+$] and the transversality of the 1-dimensional moduli spaces, together with the nature of gluing, it follows that the compactification

$$\overline{\mathcal{M}}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v})/\mathbb{R}$$

is diffeomorphic to a 1-dimensional manifold with boundary, such that

$$\partial\overline{\mathcal{M}}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v})/\mathbb{R} = \overline{\mathcal{M}}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v})/\mathbb{R} \setminus \mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(L, S, \mathbf{v})/\mathbb{R}.$$

Moreover, the boundary consists of holomorphic buildings of exactly two levels, where the upper level consists of one pseudo-holomorphic disk in

$$\mathcal{M}^{\delta,\mathbf{k}'}_{a;\mathbf{b}',\mathbf{w}';A'}(L, S, \mathbf{v}),$$

and where the lower level consists of one pseudo-holomorphic disk in

$$\mathcal{M}^{\delta,\mathbf{k}''}_{b'_i;\mathbf{b}'',\mathbf{w}'';A''}(L, S, \mathbf{v}),$$

together with trivial strips over Reeb chords, and where these moduli spaces satisfy

$$\begin{cases} \dim \mathcal{M}^{\delta,\mathbf{k}'}_{a;\mathbf{b}',\mathbf{w}';A'}(L,S,\mathbf{v}) = \dim \mathcal{M}^{\delta,\mathbf{k}''}_{b'_i;\mathbf{b}'',\mathbf{w}'';A''}(L,S,\mathbf{v}) = 1, \\ A' + A'' = A, \\ \mathbf{b} = b'_1 \dots b'_{i-1} b''_1 \dots b''_{m''} b'_{i+2} \dots b'_{m'}, \\ \mathbf{w} = (w'_1, \dots, w'_i + w''_1, w''_2, \dots, w''_{m''}, w''_{m''+1} + w'_{i+1}, \dots, w'_{m'+1}), \\ d' = w'_1 + \dots + w'_{m'+1}, \\ d'' = w''_1 + \dots + w''_{m''+1}, \\ \mathbf{k}' = (0, \dots, w'_1 + \dots + w'_i - 1, w'_1 + \dots + w'_i + d'', \dots, d' + d'' - 1), \\ \mathbf{k}'' = (w'_1 + \dots + w'_i, \dots, w'_1 + \dots + w'_i + d'' - 1). \end{cases}$$

Conversely, gluing two such solutions produces a solution inside the 2-dimensional moduli space

$$\mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v}).$$

By Proposition 9.4 it follows that, for $\delta > 0$ small enough, there is a bijection between the boundary-points $\partial\overline{\mathcal{M}}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(L,S,\mathbf{v})$ and the disjoint unions of the spaces

$$\mathcal{M}_{a;\mathbf{b}',\mathbf{w}';A'}(L,S,\mathbf{v}) \times \mathcal{M}_{b'_i;\mathbf{b}'',\mathbf{w}'';A''}(L,S,\mathbf{v})$$

satisfying

$$\begin{cases} \dim \mathcal{M}_{a;\mathbf{b}',\mathbf{w}';A'}(L,S,\mathbf{v}) = \dim \mathcal{M}_{b'_i;\mathbf{b}'',\mathbf{w}'';A''}(L,S,\mathbf{v}) = 1, \\ A' + A'' = A, \\ \mathbf{b} = b'_1 \dots b'_{i-1} b''_1 \dots b''_{m''} b'_{i+2} \dots b'_{m'}, \\ \mathbf{w} = (w'_1, \dots, w'_i + w''_1, w''_2, \dots, w''_{m''}, w''_{m''+1} + w'_{i+1}, \dots, w'_{m'+1}). \end{cases}$$

This implies that the coefficient of the word $\mathbf{b}$ in the expression ${\partial_{S,N}}^2(a)$ is the cardinality of the boundary points of a 1-dimensional manifold, and hence it is equal to 0 modulo 2.

It now remains to show that

$$(\mathcal{A}(L), \partial) \simeq (\mathcal{A}(L,S)/\langle s \rangle, \partial_{S,N}).$$

To that end, observe that a 1-dimensional moduli space $\mathcal{M}_{a;\mathbf{b},0;A}(V,S,\mathbf{v})$ is the usual moduli spaces in the definition of the Legendrian contact homology DGA. It follows that the differential $\partial_{S,N}$ is on the form

$$\partial_{S,N}(a) = \partial(a) + h_{S,N}(a), \;\; a \in \mathcal{Q}(L)_E$$

when evaluated on a generator, where ∂ is the differential for the Legendrian contact homology DGA of L and where the function $h_{S,N}$ has image in the two-sided ideal $\langle s \rangle$. $\square$

5. The isomorphism

Let Y be a contact manifold of dimension $2n+1$ and consider $L_S \subset Y$ obtained from the Legendrian submanifold $L \subset Y$ by a Legendrian ambient surgery along the k-sphere $S \subset L$ with bounding disk $D_S \subset Y$. Let $V_\epsilon \subset \mathbb{R} \times Y$ be the exact Lagrangian cobordism from L to L_S constructed in Section 3.4 corresponding to a $(k+1)$-handle attachment. Here we assume that $k < n-1$.

Fix any number $E > 0$. In this section we will only consider the Reeb chords on L and L_S having action less than E. Thus, there are finitely many such Reeb chords and they all are at some positive distance away from S. Since D_S is subcritical, we may assume that they all are disjoint from D_S as well. It follows that, after choosing $\epsilon > 0$ small enough, the Reeb chords on L_S of action less than E coincide with the Reeb chords on L of action less than E together with the Reeb chord c_{new}. We refer to Section 6 for a direct limit construction which takes care of the general situation.

As described in Section 2.5, the exact Lagrangian cobordism V_ϵ induces a DGA morphism from $(\mathcal{A}(L_S), \partial)$ to $(\mathcal{A}(L), \partial)$ by counting rigid disks in the cobordism. We will modify the definition of this chain map by taking into account rigid disks having incidence conditions on parallel copies of the core disk of the $(k+1)$-handle of the cobordism V_ϵ. This will map will be a *tame isomorphism* of DGAs

$$\Psi\colon (\mathcal{A}(L_S), \partial) \to (\mathcal{A}(L, S), \partial_{S,N}),$$

and, in particular, it will induce an isomorphism of the corresponding homologies.

5.1. **The relevant moduli spaces.** Recall the definition of the model of an exact Lagrangian $(k+1)$-handle attachment

$$W_\epsilon \subset T^*\mathbb{R}^{n+1} \simeq \mathbb{R} \times J^1(\mathbb{R}^n)$$

defined in Section 3.3. This is an exact Lagrangian cobordism from the Legendrian submanifold $L_{0,k} \subset J^1(\mathbb{R}^n)$ to $L_{\epsilon,k} \subset J^1(\mathbb{R}^n)$, where the latter is obtained by a Legendrian ambient surgery on $L_{0,k}$ along the k-sphere $S_{0,k} \subset L_{0,k}$ with bounding disk $D_{0,k} \subset J^1(\mathbb{R}^n)$.

Recall the definition of φ_ϵ in Section 3.3 and consider the submanifold

$$\widetilde{S}_{0,k} := W_\epsilon \cap \left\{ \begin{array}{l} x_1^2 + \ldots + x_{k+1}^2 = \eta - \varphi_\epsilon(\mathbf{x}), \\ x_{k+2} = \ldots = x_n = 0 \end{array} \right\} \subset W_\epsilon.$$

This submanifold may be considered as the core disk of the $(k+1)$-handle of the cobordism W_ϵ. Observe that the submanifold of $\mathbb{R} \times J^1(\mathbb{R}^n) \simeq T^*\mathbb{R}^{n+1}$ identified with $\widetilde{S}_{0,k}$ coincides with $(-\infty, -1) \times S_{0,k}$ outside of a compact set.

By $\widetilde{\mathbf{v}}_0/x_{n+1}$ we denote the vector field

$$x_1\left(1 + \frac{\partial \varphi_\epsilon}{\partial x_1}(\mathbf{x})\right)\partial_{y_1} + \ldots + x_{k+1}\left(1 + \frac{\partial \varphi_\epsilon}{\partial x_{k+1}}(\mathbf{x})\right)\partial_{y_{k+1}} + \frac{\partial \varphi_\epsilon}{\partial x_{n+1}}(\mathbf{x})\partial_{y_{n+1}}$$

along $\widetilde{S}_{0,k}$. This section can be extended to a framing

$$\widetilde{\mathbf{v}}_0, \partial_{x_{k+2}}, \ldots, \partial_{x_n},$$

of the normal bundle of $\widetilde{S}_{0,k} \subset W_\epsilon$.

Observe that the vector field $\widetilde{\mathbf{v}}_0$ restricted to

$$(-\infty, -1) \times S_{0,k} \subset \widetilde{S}_{0,k}$$

is $\mathbb{R}$-invariant and normal to

$$(-\infty, -1) \times S_{0,k} \subset (-\infty, -1) \times L_{0,k}.$$

By the construction of V_ϵ there is an exact symplectomorphism which identifies a neighborhood of $\mathbb{R} \times D_S \subset \mathbb{R} \times Y$ with the neighborhood $U_\epsilon \subset T^*(\mathbb{R}^{n+1}) \simeq \mathbb{R} \times J^1(\mathbb{R})$ defined in Section 3.4.2 such that $W_\epsilon \cap U_\epsilon \subset T^*(\mathbb{R}^{n+1})$ is identified with V_ϵ. Moreover, this exact symplectomorphism preserves the symplectization coordinate.

We will use $\widetilde{S} \subset V_\epsilon$ to denote the submanifold identified with $\widetilde{S}_{0,k} \subset T^*(\mathbb{R}^{n+1})$ under this identification, and $\widetilde{\mathbf{v}}$ to denote the vector field identified with $\widetilde{\mathbf{v}}_0$.

It follows that $\widetilde{S} \subset V_\epsilon$ coincides with $(-\infty, -1) \times S$ outside of a compact subset and that the vector field $\widetilde{\mathbf{v}}$, which is a normal to $\widetilde{S} \subset V_\epsilon$, is $\mathbb{R}$-invariant when restricted to $(-\infty, -1) \times S$. We let $\mathbf{v}$ denote the extension of $\widetilde{\mathbf{v}}|_{(-\infty,-1)\times S}$ to an $\mathbb{R}$-invariant vector field along $\mathbb{R} \times S$.

Let $\delta > 0$, $a \in \mathcal{Q}(L_S)_E$, $\mathbf{b} = b_1 \cdot \ldots \cdot b_m$ a word of Reeb chords on L, $\mathbf{w} = (w_1, \ldots, w_{m+1}) \in (\mathbb{Z}_{\geq 0})^{m+1}$ a tuple of integers, $\mathbf{k} = (k_1, \ldots, k_{w_1+\ldots+w_{m+1}}) \in (\mathbb{Z}_{\geq 0})^{w_1+\ldots+w_{m+1}}$ a tuple of increasing integers, and $A \in H_1(V)$. In Section 9 we define the moduli spaces

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon, \widetilde{S}, \widetilde{\mathbf{v}})$$

which count pseudo-holomorphic disks $u \colon \dot{D} \to \mathbb{R} \times Y$, for any choice of conformal structure on $\dot{D} = D \setminus \{p_0, \ldots, p_m\}$, satisfying

- $u(\partial D^2) \subset V_\epsilon$ and, after closing up the boundary with appropriately oriented capping paths, one obtains $A \in H_1(V_\epsilon)$.
- u has a positive puncture asymptotic to a at p_0.
- u has negative punctures asymptotic to b_i at p_i.
- u maps a number w_i of boundary points on the arc between p_i and p_{i+1} (where we set $p_{m+1} := p_0$) to the parallel copies $\exp_{\widetilde{S}}(k_i \delta \widetilde{\mathbf{v}})$ of $\widetilde{S}$, such that the i:th boundary point with respect to the order on $\partial D \setminus \{p_0\}$ induced by the orientation is mapped to $\exp_{\widetilde{S}}(k_i \delta \widetilde{\mathbf{v}})$. Here exp denotes the exponential map on V_ϵ for some choice of metric.

In the case when $\mathbf{k} = (0, 1, \ldots, w_1 + \ldots + w_{m+1} - 1)$, we will use the notation

$$\mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(V, \widetilde{S}, \widetilde{\mathbf{v}}) := \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V, \widetilde{S}, \widetilde{\mathbf{v}}).$$

Lemma 5.1. *There is a cylindrical almost complex structure on* $\mathbb{R} \times Y$*, which is integrable in a neighborhood of* $\mathbb{R} \times D_S$*, for which the moduli spaces*

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon, \widetilde{S}, \widetilde{\mathbf{v}})$$

and

$$\mathcal{M}_{a;\mathbf{b};A}(\mathbb{R} \times L_S)$$

are transversely cut out. Moreover, the almost complex structure can be chosen such that Lemma 4.2 is fulfilled, i.e. such that the moduli spaces

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R} \times L, \mathbb{R} \times S, \mathbf{v})$$

of expected dimension 1, and all moduli spaces

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R} \times L, \mathbb{R} \times S, \mathbf{v}),$$

are transversely cut out.

Proof. In the neighborhood of $\mathbb{R} \times D_S$ which is identified with $U_\epsilon \subset T^*\mathbb{R}^{n+1}$ by the above identification, we define the cylindrical almost complex structure by requiring the projections

$$(x_i, y_i/x_{n+1})\colon\ U_\epsilon \to \mathbb{C}$$

to be holomorphic for $i = 1, \ldots, n$. As noted in the proof of Lemma 4.2, this requirement makes the almost complex structure integrable in U_ϵ

Observe that we can assume that the almost complex structure constructed here coincides with the one constructed in the proof of Lemma 4.2, and hence that the statements in that lemma holds for this choice of cylindrical almost complex structure as well.

By using Proposition 9.1, we get transversality for every moduli spaces in the statement with an almost complex structure as above except those having a positive puncture at c_{new}. Namely, for these moduli spaces, Proposition 9.1 would require a perturbation of the almost complex structure in the neighborhood U_ϵ, which we want to avoid. However, the transversality for these moduli spaces follows by Lemma 5.6.

To that end, for achieving transversality of the disks with a positive puncture at c_{new} without perturbing the complex structure near $\mathbb{R} \times c_{new}$, we must instead perturb the vector field $\widetilde{\mathbf{v}}$.

Observe that the submanifold

$$W_\epsilon \cap \left\{x_1^2 + \ldots + x_{k+1}^2 = 0\right\} \subset T^*(\mathbb{R}^{n+1}),$$

which may be viewed as the cocore disk of the $(k+1)$-handle attachment W_ϵ, intersects the core disk $\widetilde{S}_{0,k}$ transversely precisely at the point in $T^*\mathbb{R}^{n+1}$ satisfying

$$\begin{cases} x_i = 0, & i \leq n, \\ y_i = 0, & i \leq n+1, \\ x_{n+1} = 1. \end{cases}$$

Moreover, the vector field $\widetilde{\mathbf{v}}_0$ is tangent to the cocore disk at this point. A perturbation of $\widetilde{\mathbf{v}}_0$ in a neighborhood of this point will take us out of this degenerate situation. □

5.2. **The tame isomorphism Ψ.** Let $(\mathcal{A}(L_S)^E, \partial) \subset (\mathcal{A}(L_S), \partial)$ denote the sub-complex generated by Reeb chords of action less than E.

We define the map

$$\Psi\colon\ \mathcal{A}(L_S)^E \to \mathcal{A}(L,S)^E$$

on the generators $\mathcal{Q}(L_S)_E$ of $\mathcal{A}(L_S)^E$ by

$$\Psi(a) = \sum_{\dim \mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}})=0} |\mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}})| s^{w_1} b_1 s^{w_2} \cdot \ldots \cdot s^{w_m} b_m s^{w_{m+1}}.$$

The map Ψ is then extended to all of $\mathcal{A}(L_S)^E$ as an algebra map.

Observe that by Formula (5.1) it follows that if $b_1 \cdot \ldots \cdot b_m$ is a word of Reeb chords on L with non-zero coefficient in $\Psi(a)$, where $a \in \mathcal{Q}(L)_E$, then

$$\frac{1+2\epsilon^{1/3}}{1-2\epsilon^{1/3}} E \geq \frac{1+2\epsilon^{1/3}}{1-2\epsilon^{1/3}} \ell(a) \geq \ell(b_1) + \ldots + \ell(b_m).$$

Since the action-spectrum of the Reeb chords on L is discrete, we may thus assume that Ψ does not increase action after choosing a small enough $\epsilon > 0$ (depending on E). Furthermore, by the compactness results in Proposition 9.2, it follows that each rigid moduli space occurring the above expression is finite. Finally, by the compactness theorem in [BEH$^+$], it follows that the moduli space $\mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}})$, where a and $\mathbf{b}$ are fixed, is non-empty for only finitely many values of $A \in H_1(V)$. Consequently, Ψ is well-defined.

The dimension formula

$$\dim \mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}}) = |a| - |b_1| - \ldots - |b_m| - |s|(w_1 + \ldots + w_{m+1}) + \mu(A),$$

shows that Ψ is of degree 0. However, we have to choose the grading in the group

$$\mathbb{Z}/\mu(H_1(V_\epsilon))$$

for both $\mathcal{A}(L)$ and $\mathcal{A}(L,S)$. We refer to Section 3.5 for a discussion of the effect of a Legendrian ambient surgery on the Maslov class.

Proof of Theorem 1.4. First, the chain-map property can be proven by an argument similar to the one showing that ${\partial_{S,N}}^2 = 0$ in the proof of Theorem 1.2. To that end observe that, for $\delta > 0$ small enough, Proposition 9.2 implies that

$$|\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}})| \equiv |\mathcal{M}^{\delta,\mathbf{k}'}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}})| \pmod 2$$

for different choices of tuples $\mathbf{k}$ and $\mathbf{k}'$ whenever

$$\dim \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}}) = 0.$$

We now show that Ψ is a tame isomorphism. To that end, we may have to choose a smaller ϵ. For technical reasons, however, we want to keep the almost complex structure fixed inside the above neighborhood U_ϵ. For ϵ' satisfying $0 < \epsilon' \leq \epsilon$ we proceed as follows in order to construct $V_{\epsilon'}$. Recall that, by construction, there is a neighborhood of $\mathbb{R} \times D_S$ which is symplectically identified with $U_\epsilon \subset T^*\mathbb{R}^{n+1}$ under which V_ϵ is identified with W_ϵ. We construct $V_{\epsilon'}$ by replacing $W_\epsilon \cap U_\epsilon$ with $W_{\epsilon'} \cap U_\epsilon$, and using the above identification. Recall that Lemma 3.3 implies that W_ϵ and $W_{\epsilon'}$ coincide outside of $U_{(2/3)\epsilon}$.

After replacing V_ϵ with $V_{\epsilon'}$ as above, where $\epsilon' > 0$ is small enough, Lemma 5.5, 5.7, and Formula (5.1) imply that Ψ acts as follows on generators:

$$\begin{aligned}&\Psi(c) = c + \mathbf{b}_c, \ \ c \in \mathcal{Q}(L_S) \setminus \{c_{new}\},\\ &\Psi(c_{new}) = s,\end{aligned}$$

where $\mathbf{b}_c$ is an element in the sub-algebra generated by the Reeb chords of L different form c.

Finally, to see that $\Phi \circ \Psi^{-1}$ is the quotient projection as claimed, recall that

$$(\mathcal{A}(L)^E, \partial) \simeq (\mathcal{A}(L,S)^E, \partial_{S,N})/\langle s \rangle$$

by Theorem 1.2 and that the map Ψ is on the form $\Phi + f_S$ on generators, where f_S is a map whose image is contained in the two-sided ideal generated by s. This follows since the moduli spaces $\mathcal{M}^\delta_{a;\mathbf{b},0;A}(V_\epsilon, \widetilde{S}, \widetilde{\mathbf{v}})$ are the moduli spaces used in the definition of Φ. □

Proof of Corollary 1.5. To see the bijection of augmentations, we proceed as follows. Let I to be two-sided ideal generated by s. An augmentation of

$$(\mathcal{A}(L), \partial) \simeq (\mathcal{A}(L,S), \partial_{S,N})/I$$

pulls back to an augmentation of $(\mathcal{A}(L,S), \partial_{S,N})$. Conversely, an augmentation of $(\mathcal{A}(L,S), \partial_{S,N})$ has to map $s \mapsto 0$ since $|s| \neq 0$ holds by assumption (in the group where the grading is taken), and hence it descends to an augmentation of $(\mathcal{A}(L), \partial)$. □

5.3. Pseudoholomorphic disks in the cobordism.

We let J be a cylindrical almost complex structure on $\mathbb{R} \times Y$ as constructed in the proof of Lemma 5.1. Hence, by definition, there is a holomorphic identification of a neighborhood of $\mathbb{R} \times D_S \subset \mathbb{R} \times Y$ with the neighborhood $U_\epsilon \subset T^*\mathbb{R}^{n+1}$ as defined in Section 3.4.2, where the latter is endowed with the unique cylindrical complex structure under which the projections

$$(x_i, y_i/x_{n+1})\colon\ U_\epsilon \to \mathbb{C}$$

are holomorphic.

By assumption, we have constructed the cobordism V_ϵ. Let ϵ' be a number satisfying $0 < \epsilon' \leq \epsilon$. We construct the exact Lagrangian cobordism $V_{\epsilon'}$ by replacing $W_\epsilon \cap U_\epsilon$ with $W_{\epsilon'} \cap U_\epsilon$ inside the neighborhood of $\mathbb{R} \times D_S$ identified with U_ϵ as above. Recall that W_ϵ and $W_{\epsilon'}$ coincide outside of $U_{(2/3)\epsilon}$ according to Lemma 3.3.

Recall that the non-cylindrical part of $V_{\epsilon'}$ is contained inside the subset

$$(\log(1-(\epsilon')^{1/3}), \log(1+(\epsilon')^{1/3})) \times Y \subset \mathbb{R} \times Y$$

of the symplectization. We let

$$N_- := \log(1-2\epsilon^{1/3}), \ \ N_+ := \log(1+2\epsilon^{1/3}).$$

Suppose that D be a pseudo-holomorphic disk in $\mathbb{R} \times Y$ with boundary on $V_{\epsilon'}$ and boundary punctures asymptotic to Reeb chords of L and L_S. In the following, we fix the definition of the λ-energy of D to be

$$E_\lambda(D) := \sup_{\substack{\rho\colon (-\infty,N_-]\to\mathbb{R}_{\geq 0} \\ \int_{(-\infty,N_-)}\rho(t)dt=1}} \int_D \rho(t)dt \wedge \lambda + \sup_{\substack{\rho\colon [N_+,+\infty)\to\mathbb{R}_{\geq 0} \\ \int_{[N_+,+\infty)}\rho(t)dt=1}} \int_D \rho(t)dt \wedge \lambda,$$

and the total energy of D to be

$$E(D) = E_{d\lambda}(D) + E_\lambda(D) + \int_{D\cap([N_-,N_+]\times Y)} dt \wedge \lambda.$$

Lemma 5.2. *Let D be a punctured pseudo-holomorphic disk in the symplectization having boundary on $V_{\epsilon'}$, positive boundary punctures asymptotic to the Reeb chords*

$$c_1^+, \ldots, c_{m_+}^+ \in \mathcal{Q}(L_S),$$

and negative boundary punctures asymptotic to the Reeb chords

$$c_1^-, \ldots, c_{m_-}^- \in \mathcal{Q}(L).$$

The energies of D satisfy the following bounds:

$$0 \ \leq \ E_{d\lambda}(D) \leq \frac{1+2(\epsilon')^{1/3}}{1-2(\epsilon')^{1/3}} \sum_{i=1}^{m_+} \ell(c_i^+) - \sum_{i=1}^{m_-} \ell(c_i^-), \tag{5.1}$$

$$0 \ \leq \ E_\lambda(D) + \int_{D\cap([N_-,N_+]\times Y)} dt \wedge \lambda \leq \frac{2}{1-2\epsilon^{1/3}} \ell(c_i^+). \tag{5.2}$$

Proof. The statements more or less follow from [Ekh2, Lemma B.3]. Below follows an alternative computation.

Let $\varphi\colon \mathbb{R} \to \mathbb{R}$ be a smooth function satisfying $\varphi' \geq 0$ and

$$\varphi(t) = \begin{cases} \log(1-2(\epsilon')^{1/3}), & t \leq \log(1-2(\epsilon')^{1/3}) \\ t, & \log(1-(\epsilon')^{1/3}) \leq t \leq \log(1+(\epsilon')^{1/3}) \\ \log(1+2(\epsilon')^{1/3}), & t \geq \log(1+2(\epsilon')^{1/3}). \end{cases}$$

For the bound of the $d\lambda$-energy, when $\epsilon' > 0$ is small enough, consider the inequality

$$0 \leq (1-2(\epsilon')^{1/3})E_{d\lambda}(D) = (1-2(\epsilon')^{1/3}) \int_D d\lambda \leq \int_D d(e^{\varphi(t)}\lambda)$$

for the $d\lambda$-area of pseudo-holomorphic disks D in $Y \times \mathbb{R}$.

Observe that the pullback of $e^{\varphi(t)}\lambda$ to $V_{\epsilon'}$ equals the pullback of $e^t\lambda$, since both pullbacks vanish where $V_{\epsilon'}$ is cylindrical over a Legendrian submanifold in Y. Furthermore, by construction, the primitive of the pullback of $e^t\lambda$ to $V_{\epsilon'}$ may be taken to vanish outside of a compact set.

Because of the asymptotical behavior of D, we now conclude that

$$\begin{aligned}0 &\leq (1-2(\epsilon')^{1/3})\int_D d\lambda \leq \int_D d(e^{\varphi(t)}\lambda)\\ &= \int_{\partial D} e^{\varphi(t)}\lambda\\ &= e^{\log(1+2(\epsilon')^{1/3})}\sum_{i=1}^{m_+}\ell(c_i^+) - e^{\log(1-2(\epsilon')^{1/3})}\sum_{i=1}^{m_-}\ell(c_i^-).\end{aligned}$$

For Formula (5.2) follows from the following computation. Let $\sigma : \mathbb{R} \to \mathbb{R}_{\geq 0}$ be any smooth function satisfying the following conditions:

- $\int_{(-\infty,\log(1-2(\epsilon)^{1/3})]}\sigma(t) = 1$,
- $\int_{[\log(1+2(\epsilon)^{1/3}),+\infty)}\sigma(t) = 1$,
- $\sigma(t) = \frac{1}{1-2(\epsilon)^{1/3}}e^t \geq 1$ for $t \in [\log(1-2(\epsilon)^{1/3}), \log(1+2(\epsilon)^{1/3})]$.

Let $\Sigma(t) := \int_{-\infty}^t \sigma(s)ds$. The J-holomorphicity of the disk implies the inequality

$$\begin{aligned}0 &\leq \int_D \sigma(t)dt\wedge\lambda = \int_D d(\Sigma(t)\lambda) - \int_D \Sigma(t)d\lambda\\ &\leq \int_D d(\Sigma(t)\lambda).\end{aligned}$$

Finally, we have the equality

$$\begin{aligned}\int_D d(\Sigma(t)\lambda) &= \left(\int_{\mathbb{R}}\sigma(s)ds\right)\left(\sum_{i=1}^{m_+}\ell(c_i^+)\right) + \int_{\partial D}\Sigma(t)\lambda\\ &= \frac{2}{1-2\epsilon^{1/3}}\sum_{i=1}^{m_+}\ell(c_i^+),\end{aligned}$$

where we have used that

$$\int_{\mathbb{R}}\sigma(s)ds = 2 + \frac{(1+2\epsilon^{1/3})-(1-2\epsilon^{1/3})}{1-2\epsilon^{1/3}} = 2 + \frac{4\epsilon^{1/3}}{1-2\epsilon^{1/3}} = \frac{2}{1-2\epsilon^{1/3}}$$

together with the fact that $\Sigma(t)\lambda$ is exact on V_ϵ, where the primitive can be taken to have compact support. □

Lemma 5.3. *Let C be a J-holomorphic disk in $\mathbb{R}\times Y$ with boundary on $V_{\epsilon'}$, where $0<\epsilon'\leq\epsilon$, and boundary punctures asymptotic to Reeb chords. Identify $U_\epsilon \subset T^*\mathbb{R}^{n+1}$ with a neighborhood of $\mathbb{R}\times D_S \subset \mathbb{R}\times Y$ as above. There is a constant $A>0$ independent of $\epsilon'>0$ for which the following holds. If C passes through the set identified with $\partial U_{(3/4)\epsilon} \subset U_\epsilon$, then*

$$A \leq E(C).$$

Proof. Recall that $V_{\epsilon'} \subset \mathbb{R} \times Y$ coincides with $W_{\epsilon'} \subset T^*\mathbb{R}^{n+1}$ under the identification above. Moreover, according to Lemma 3.3, $V_{\epsilon'}$ coincides with the cylindrical Lagrangian cobordism $\mathbb{R} \times L$ outside of the set identified with $U_{(2/3)\epsilon} \subset U_\epsilon$.

Consider the symplectic form

$$\omega_0 = dt \wedge \lambda + d\lambda$$

on $\mathbb{R} \times Y$ which is compatible with J and invariant under translations of the t-coordinate.

Using the identification of $T^*\mathbb{R}^{n+1}$ with $\mathbb{R} \times J^1(\mathbb{R}^n)$ as in Section 3.3, where the latter space is equipped with coordinates $(t, (\mathbf{q}, \mathbf{p}, z))$, U_ϵ is identified with the neighborhood

$$\widetilde{U}_\epsilon := \left\{ \begin{array}{l} q_1^2 + \ldots + q_{k+1}^2 < \eta + \epsilon, \\ q_{k+2}^2 + \ldots + q_n^2 < 2\epsilon, \\ t > 0 \\ |p_i| < 3\sqrt{\epsilon(\eta + \epsilon)}, \ \ i \leq n, \\ |z| < 3\epsilon^{1/6}(\eta + \epsilon) \end{array} \right\} \subset \mathbb{R} \times J^1(\mathbb{R}^n),$$

which is invariant under translations of the t-coordinate, while the above symplectic form is identified with the symplectic form

$$dt \wedge \left(dz - \sum_i p_i dq_i \right) + \sum_i dq_i \wedge dp_i.$$

The monotonicity property for pseudo-holomorphic curves with and without boundary (see [Sik, Proposition 7.3.1] and [Sik, Proposition 7.4.3]), respectively, now applies to pseudo-holomorphic curves as above passing through $\partial U_{(3/4)\epsilon} \subset \mathbb{R} \times J^1(\mathbb{R}^n)$.

In particular, there is a covering of $\partial \widetilde{U}_{(3/4)\epsilon}$ by a t-invariant set of open balls of radius r (relative the above coordinates) contained inside $\widetilde{U}_\epsilon$, where $r_0 < r < 1$ for some fixed $r_0 > 0$. The above monotonicity property provides a constant $A > 0$ with the property that any disk C having either a boundary-point or an internal point intersecting $\partial \widetilde{U}_{(3/4)\epsilon}$ has ω_0-area bounded from below by A when intersected with one of the above balls. The same estimate on the total energy of such a disk C now follows, which finishes the proof. □

Lemma 5.4. *For $\epsilon' > 0$ small enough every J-holomorphic disk D in the symplectization which has boundary on $V_{\epsilon'}$ and exactly one puncture, which moreover is positive and asymptotic to c_{new}, is contained in a set identified with*

$$U_\epsilon \cap \left\{ \begin{array}{l} x_1 = \ldots = x_{k+1} = y_1 = \ldots = y_{k+1} = 0, \\ x_{n+1} \geq 1 \end{array} \right\} \subset T^*\mathbb{R}^{n+1}$$

under the above identification.

Proof. Any disk satisfying the assumptions of the lemma, but that is not contained in U_ϵ, must intersect $\partial U_{(3/4)\epsilon}$ and hence has total energy bounded by A from below by Lemma 5.3.

Formulas (5.1) and (5.2) show that for $\epsilon' > 0$ small enough, we may assume that the total energy of D is arbitrary small. Together with the above monotonicity property, we may hence assume that D is contained inside U_ϵ for $\epsilon' > 0$ small enough.

Consider the holomorphic projections

$$(x_i, y_i/x_{n+1})\colon\ U_\epsilon \to \mathbb{C}$$

for $i = 1, \ldots, n$. The image of

$$W_{\epsilon'} \cap U_\epsilon$$

under these projections is shown in Figure 1 for $i = 1, \ldots, k+1$, and in Figure 2 for $i = k+2, \ldots, n$. We use f_i to denote the composition of the disk D with the projection $(x_i, y_i/x_{n+1})$.

By the asymptotical property of D we know that each f_i fills some part of the corner above or below the image of c_{new} in Figure 1 for $i = 1, \ldots, k+1$, unless f_i is constantly zero. By the open-mapping theorem it thus follows that $f_i = 0$.

Furthermore, considering the projections f_i for $i = k+2, \ldots, n$, the open mapping theorem implies that

$$D \subset U_\epsilon \cap \{x_1 = \ldots = x_{k+1} = y_1 = \ldots = y_{k+1} = 0\}.$$

Finally, to see that $x_{n+1} \geq 1$ on D, observe that this estimate is true for the boundary of D since

$$W_{\epsilon'} \cap \{x_1 = \ldots = x_{k+1} = y_1 = \ldots = y_{k+1} = 0\} \subset \{x_{n+1} \geq 1\}.$$

The result now follows by the open mapping theorem applied to the projection to the (x_{n+1}, y_{n+1})-plane restricted to D. □



FIGURE 1. The image of $W_{\epsilon'}$ under the holomorphic projections $(x_i, y_i/x_{n+1})$ with $i = 1, \ldots, k+1$.



FIGURE 2. The image of $W_{\epsilon'}$ under the holomorphic projections $(x_i, y_i/x_{n+1})$ with $i = k+2, \dots, n$.

Lemma 5.5. *For small enough $\epsilon' > 0$, every J-holomorphic disk D in the symplectization which has boundary on $V_{\epsilon'}$ intersecting $\widetilde{S}$ and exactly one puncture, which moreover is positive and asymptotic to c_{new}, is contained in the holomorphic strip $\mathbb{R} \times c_{new} \subset \mathbb{R} \times Y$. In particular, there is a unique such disk.*

Proof. Let D be such a disk. By the previous lemma, we may assume that

$$D \subset U_\epsilon \cap \{x_1 = \ldots = x_{k+1} = y_1 = \ldots = y_{k+1} = 0,\ x_{n+1} \geq 1\}$$

in $T^*\mathbb{R}^{n+1}$. Thus, D intersects $\widetilde{S}$ in the set

$$\widetilde{S}_{0,k} \cap U_\epsilon \cap \{x_1 = \ldots = x_{k+1} = 0\} = \left\{ \begin{array}{l} x_i = y_i = 0, \quad i = 1, \ldots, n, \\ x_{n+1} = 1, \\ y_{n+1} = 0 \end{array} \right\}.$$

The holomorphic projections

$$(x_i, y_i/x_{n+1})\colon\ U_\epsilon \to \mathbb{C},$$

for $i = k+2, \ldots, n$ maps

$$W_{\epsilon'} \cap \{x_1 = \ldots = x_{k+1} = 0,\ \ x_{n+1} \geq 1\}$$

to a filled figure-8 having its center at the origin in $\mathbb{C}$ as shown in Figure 2. Let f_i denote the composition of D with the projection $(x_i, y_i/x_{n+1})$, which is holomorphic on the disk.

By using the open mapping theorem, we get that the image of f_i restricted d to D has values in the filled figure-8 and, moreover, if a boundary point of D is mapped to the origin in Figure 2 by f_i, it must vanish identically. We conclude that D is contained in the holomorphic strip $\mathbb{R} \times c_{new}$. □

Lemma 5.6. *The moduli spaces $\mathcal{M}_{c_{new};0,\mathbf{w};A}(V_\epsilon, \widetilde{S}, \widetilde{\mathbf{v}})$ with $\mathbf{w} \neq 0$, as well as the moduli space $\mathcal{M}_{c_{new};0;A}(L_S)$ in the case $|c_{new}| = 1$, are all transversely cut out for this particular choice of J.*

Proof. We start with the spaces $\mathcal{M}_{c_{new};0,\mathbf{w};A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}})$, where $\mathbf{w}\neq 0$. Observe that by Lemma 5.5, together with the choice of $\widetilde{\mathbf{v}}$ in the proof of Lemma 5.1, in order for the moduli space to be non-empty, we must have $\mathbf{w}=1$. Moreover, there is a unique disk in this moduli space.

Since the disk is contained in U_ϵ, the linearized problem splits into the summands

$$(E_1,F_1)\oplus(E_2,F_2)\oplus(E_3,F_3),$$

where $(E_i,F_i)\to D^2\setminus\{1\}$ are complex bundles with a real sub-bundle $F_i\to\partial D^2\setminus\{1\}$ over the punctured unit disk. Here, we let E_1 correspond to the coordinates $(x_1,\ldots,x_{k+1},y_1,\ldots,y_{k+1})$, E_2 correspond to the coordinates $(x_{k+2},\ldots,x_n,y_{k+2},\ldots,y_n)$ and E_3 correspond to the coordinates (x_{n+1},y_{n+1}). We have natural holomorphic trivializations

$$E_1\simeq\mathbb{C}^{k+1},\;\;E_2\simeq\mathbb{C}^{n-k-1},\;\;E_3\simeq\mathbb{C},$$

under which

$$\begin{aligned}F_1&\simeq\mathbb{R}\langle e^{i\varphi_1(\theta)}\mathbf{e}_j;\;\;1\leq j\leq k+1\rangle\subset E_1,\\F_2&\simeq\mathbb{R}\langle e^{i\varphi_2(\theta)}\mathbf{e}_j;\;\;k+2\leq j\leq n\rangle\subset E_2,\\F_3&\simeq\mathbb{R}e^{i\varphi_3(\theta)}\mathbf{e}_{n+1}\subset E_3,\end{aligned}$$

where θ is a coordinate on ∂D^2.

One can check that φ_1 is decreasing and has range inside $(-\pi/4,\pi/4)$, φ_2 is increasing and has range inside $(-\pi/4,\pi/4)$, and that φ_3 is increasing and has range inside $[0,\pi]$.

Observe that since we have an integrable complex structure in a neighborhood of the disk, the linearized operator is the standard Cauchy-Riemann operator.

The adjoint problem can naturally be considered to have boundary condition

$$(E_1,F_1^*)\oplus(E_2,F_2^*)\oplus(E_3,F_3^*)\simeq((E_1,F_1)\oplus(E_2,F_2)\oplus(E_3,F_3))^*$$

given by

$$\begin{aligned}F_1^*&\simeq\mathbb{R}\langle e^{-i(\varphi_1(\theta)+\theta)}\mathbf{e}_j;\;\;1\leq j\leq k+1\rangle\subset E_1,\\F_2^*&\simeq\mathbb{R}\langle e^{-i(\varphi_2(\theta)+\theta)}\mathbf{e}_j;\;\;k+2\leq j\leq n\rangle\subset E_2,\\F_3^*&\simeq\mathbb{R}e^{-i(\varphi_3(\theta)+\theta)}\mathbf{e}_{n+1}\subset E_3.\end{aligned}$$

By the argument principle it now follows that the adjoint problem has no kernel, and hence, the original problem has no cokernel.

To check that the sections in the kernel are transverse to $T\widetilde{S}$ at the intersection-point, we observe that non-zero sections in the kernel cannot vanish at any boundary point except the puncture by the open mapping principle. This concludes the proof that $\mathcal{M}_{c_{new};0,1;A}(V_\epsilon,\widetilde{S},\widetilde{\mathbf{v}})$ is transversely cut out.

We now show the transversality for $\mathcal{M}_{c_{new};0;A}(L_S)$ when $|c_{new}|=n-k-1=1$, and hence $k=n-2$. Since D has no negative punctures, we may identify each solution D in $\mathcal{M}_{c_{new};0;A}(L_S)$ with a solution having boundary on $V_\epsilon \subset \mathbb{R} \times Y$ and satisfying $\min_t D \geq \log 2$, where t is the symplectization coordinate.

Lemma 5.4 now implies that D is contained in a neighborhood identified with

$$U_\epsilon \cap \{x_1 = \ldots = x_{n-1} = 0 = y_1 = \ldots = y_{n-1},\ x_{n+1} \geq 2\} \subset T^*\mathbb{R}^{n+1}$$

under the above exact symplectomorphism. Considering Figure 2, we immediately find that a solution has to map to either of the two domains bounded by the figure-8 curve under the projection to the $(x_n, y_n/x_{n+1})$-plane. Conversely, by using the methods in [ENS, Section 7.2], one can show that there are unique lifts of the two obvious disks in the $(x_n, y_n/x_{n+1})$-plane bounded by the figure-8 curve to disks in the symplectization having a positive puncture at c_{new}.

To see that these disks are transversely cut out, again the linearized problem splits up into the summands

$$(E_1, F_1) \oplus (E_2, F_2) \oplus (E_3, F_3).$$

In this case, the boundary condition is given by

$$\begin{aligned} F_1 &\simeq \mathbb{R}\langle e^{i\varphi_1(\theta)}\mathbf{e}_j;\ 1 \leq j \leq n-1\rangle \subset E_1, \\ F_2 &\simeq \mathbb{R}e^{i\varphi_2(\theta)}\mathbf{e}_n \subset E_2, \\ F_3 &\simeq \mathbb{R}e^{i\varphi_3(\theta)}\mathbf{e}_{n+1} \subset E_3, \end{aligned}$$

where φ_1 is decreasing and has range inside $[-\pi/4, \pi/4]$, φ_2 is increasing and has range in $[\pi/4, 7\pi/4]$ and φ_3 is increasing has range in $[0, \pi]$. Consequently, the dual of the linearized problem has no kernel, and hence the linearization of the original problem has no cokernel. □

Lemma 5.7. *Let $c \in \mathcal{Q}(L_S) \setminus \{c_{new}\}$. For $\epsilon' > 0$ small enough, the only disk in $\mathcal{M}_{c;c;A}(V_{\epsilon'})$ is the trivial strip over c.*

Proof. Recall that we have fixed an identification of a neighborhood of $\mathbb{R} \times D_S \subset \mathbb{R} \times Y$ with $U_\epsilon \subset T^*\mathbb{R}^{n+1}$. Any disk inside $\mathcal{M}_{c;c;A}(V_{\epsilon'})$ which is not equal to a trivial strip must have boundary intersecting $\partial U_{(2/3)\epsilon}$ under this identification since, otherwise, the disk would belong to $\mathcal{M}_{c;c;A}(L)$, and would hence have vanishing $d\lambda$-energy, contradicting the assumption that the disk is not a trivial strip.

We assume the contrary. Suppose that there is a sequence $\epsilon_i > 0$ satisfying $\epsilon_i \to 0$ as $i \to \infty$, such that there is a non-trivial solution D_i

$$(a_i(s,t), u_i(s,t))\colon\ \mathbb{R} \times [0,1] \to D_i \subset \mathbb{R} \times Y$$

inside $\mathcal{M}_{c;c;A}(V_{\epsilon_i})$ for each i, where s is the standard coordinate on the $\mathbb{R}$-factor, t is the standard coordinate on the $[0,1]$-factor, and where $\mathbb{R} \times [0,1] \subset \mathbb{C}$ is endowed with the standard complex structure. Observe that $E_{d\lambda}(D_i) \to 0$ as $i \to \infty$ according to Formula (5.1).

Consider the path-space $C^\infty(([0,1],\{0,1\}),(Y,L_S))$ of paths with endpoints on L_S. Fix a small enough neighborhood O of $c \subset Y$ with the property that the only Reeb chord on L_S contained inside O is c, and that $O \cap L_S = O \cap L$. Take sufficiently small neighborhoods $U_1, U_2, U_3 \subset C^\infty(([0,1],\{0,1\}),(Y,L_S))$ of the parametrized Reeb-chord c satisfying the following.

- $c \in U_1 \subset U_2 \subset U_3$,
- U_3 consists of paths contained inside O,
- $U_i = B_{i\delta}(c) \cap C^\infty(([0,1],\{0,1\}),(Y,L_S))$ for some $\delta > 0$, where $B_{i\delta}(c) \subset C^1(([0,1],\{0,1\}),(Y,L_S))$ denotes a ball of radius $i\delta$ centered at c with respect to some choice of uniform C^1-norm.

Using the asymptotical properties of J-holomorphic curves in symplectizations (see [HWZ]), and the assumption that D_i is not a trivial strip, for each solution D_i we can select some $s_i \gg 0$ for which $u(s_i,\cdot) \in U_2 \setminus U_1$ and $u(s,\cdot) \in U_2$ when $s \geq s_i$.

A standard argument implies that no bubbling can occur inside the neighborhood $\mathbb{R} \times O$. Hence, there are choices of numbers b_i for which the sequence

$$(a_i(s-s_i,t)-b_i, u_i(s-s_i,t))\colon\ [0,+\infty)\times[0,1] \to D_i \subset \mathbb{R}\times Y$$

of J-holomorphic strips converges to a J-holomorphic strip

$$(a_\infty(s,t), u_\infty(s,t))\colon\ [0,+\infty)\times[0,1] \to D_i \subset \mathbb{R}\times Y$$

of finite energy being asymptotic to c as $s \to +\infty$. It follows that $u_\infty(0,\cdot) \notin U_1$ by construction, which contradicts the fact that the $d\lambda$-energy of the limit vanishes. □

6. The case of infinitely many Reeb chords on the handle

In the case when infinitely many Reeb chords are present, one has to handle the situation by applying Theorem 1.4 restricted to the Reeb chords below each finite action level, and then taking direct limits.

We construct a sequence

$$V_{\epsilon_0}, V_{\epsilon_1}, V_{\epsilon_2}, V_{\epsilon_3}, \ldots,$$

of exact Lagrangian cobordisms, where

$$\epsilon_0 := \epsilon > \epsilon_1 > \epsilon_2 > \epsilon_3 > \ldots > 0,\ \ \epsilon_i \to 0,$$

and

$$0 < E_0 < E_1 < E_2 < E_3 < \ldots,\ \ E_i \to \infty,$$

such that each V_{ϵ_i} satisfies the results in Section 5 for the Reeb chords of action less than E_i and some choice of cylindrical almost complex structure J_i.

More precisely, we may assume that the following holds. Below the action level E_i, the Reeb chords on L_{S,ϵ_i} correspond to the Reeb chords on L together with the Reeb chord c_{new}, where we define L_{S,ϵ_i} to be the Legendrian at the positive end of V_{ϵ_i}. Moreover, there is a DGA-isomorphism

$$\Psi_i\colon (\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i) \to (\mathcal{A}(L,S)^{E_i}, \partial_{S,N,i})$$

such that the DGA-morphism

$$\Phi_i\colon (\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i) \to (\mathcal{A}(L)^{E_i}, \partial_i)$$

induced by V_{ϵ_i} satisfies the property that $\Phi_i \circ \Psi^{-1}$ corresponds to the quotient projection

$$(\mathcal{A}(L,S)^{E_i}, \partial_{S,N,i}) \to (\mathcal{A}(L,S)^{E_i}/\langle s\rangle, \partial_{S,N,i}).$$

We may assume that J_i and J_j agree in some neighborhood of the handle. Moreover, we can assume that J_i and J_{i+1} are close enough, so that

$$\partial_{S,N,i+1}|_{\mathcal{A}(L,S)^{E_i}} = \partial_{S,N,i}$$

and

$$\partial_{i+1}|_{\mathcal{A}(L)^{E_i}} = \partial_i.$$

Namely, if J_{i+1} is a small enough perturbation of J_i, we may assume that there is a bijective correspondence of the solutions which are transversely cut out before the perturbation with the corresponding solutions after the perturbation.

In this way, we obtain an almost complex structure J_∞ defining the DGA $(\mathcal{A}(L), \partial)$, which thus has a filtration

$$\ldots \subset (\mathcal{A}(L)^{E_i}, \partial_i) \subset (\mathcal{A}(L)^{E_{i+1}}, \partial_{i+1}) \subset \ldots \subset (\mathcal{A}(L), \partial).$$

There exists invertible exact Lagrangian cobordisms from $L_{S,\epsilon_{i+1}}$ to L_{S,ϵ_i} which can be constructed out of a part of

$$V_{\epsilon_i} \cap (\log(1), +\infty) \times Y.$$

These cobordisms induce DGA quasi-isomorphisms

$$\alpha_i\colon (\mathcal{A}(L_{S,\epsilon_i}), \partial_i) \to (\mathcal{A}(L_{S,\epsilon_{i+1}}), \partial_{i+1})$$

by the results in [Ekh2] (also, see Section 2.5). After passing to a subsequence of ϵ_i, and after constructing the cobordisms with some care, the induced maps α_i can be made to satisfy the following property.

Lemma 6.1. *If $\sum_{i=0}^{\infty}(\epsilon_i)^{1/3} < \infty$, then we may assume that the maps α_i above have the property that for any $x \in \mathcal{A}(L_{S,\epsilon_i})^E$ and $M \in \mathbb{Z}_{\geq 0}$,*

$$\alpha_{i+M} \circ \alpha_{i+M-1} \circ \ldots \alpha_i(x) \in \mathcal{A}(L_{S,\epsilon_{i+M+1}})^{CE},$$

where $C \geq 1$ is a constant independent of x, M and E.

Proof. We may assume that the exact Lagrangian cobordism inducing α_i is cylindrical outside of the set

$$(\log(1), \log(1 + (\epsilon_i)^{1/3})) \times Y.$$

Applying [Ekh2, Lemma B.3], or alternatively, arguing as in the derivation of Formula (5.1), it follows that a pseudo-holomorphic disk D with a positive puncture asymptotic to a and negative punctures asymptotic to $b_1, \ldots, b_m$ in the same cobordism satisfy

$$0 \leq E_{d\lambda}(D) \leq (1+2(\epsilon_i)^{1/3})\ell(a) - \sum_{i=1}^{m} \ell(b_i),$$

and consequently

$$\ell(b_i) \leq (1+2(\epsilon_i)^{1/3})\ell(a).$$

This implies that

$$\alpha_i(a) \in \mathcal{A}(L_{S,\epsilon_{i+1}})^{(1+2(\epsilon_i)^{1/3})\ell(a)}$$

holds for a generator $a \in \mathcal{A}(L_{S,\epsilon_i})$, from which the claim follows by induction. $\square$

Using the above lemma it follows that, after possibly passing to a subsequence of ϵ_i, we may suppose that the maps α_i induce DGA-morphisms

$$\alpha_i \colon (\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i) \to (\mathcal{A}(L_{S,\epsilon_{i+1}})^{E_{i+1}}, \partial_{i+1}).$$

We define $(\mathcal{A}(L_S), \partial) := (\mathcal{A}(L_{S,\epsilon_0}), \partial_0)$.

Lemma 6.2.

$$\lim_{i\to\infty} H(\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial) \simeq H(\mathcal{A}(L_S), \partial).$$

Proof. This follows by studying the commutative diagram of DGA-morphisms

$$\begin{tikzcd}
\cdots \arrow[r, "\alpha_{i-1}"] & \mathcal{A}(L_{S,\epsilon_i}) \arrow[r, "\alpha_i"] & \mathcal{A}(L_{S,\epsilon_{i+1}}) \arrow[r, "\alpha_{i+1}"] & \cdots \\
\cdots \arrow[r, "\alpha_{i-1}"] & \mathcal{A}(L_{S,\epsilon_i})^{E_i} \arrow[r, "\alpha_i"] \arrow[u, hook] & \mathcal{A}(L_{S,\epsilon_{i+1}})^{E_{i+1}} \arrow[r, "\alpha_{i+1}"] \arrow[u, hook] & \cdots,
\end{tikzcd}$$

and observing that on the homology level the above sequence consists of isomorphisms.

By taking direct limits of the two sequences, we get an inclusion

$$\lim_{i\to\infty} H(\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i) \hookrightarrow \lim_{i\to\infty} H(\mathcal{A}(L_{S,\epsilon_i}), \partial_i) \simeq H(\mathcal{A}(L_S), \partial).$$

To see that this map is surjective, we observe that every fixed element $y \in H(\mathcal{A}(L_{S,\epsilon_0}), \partial_0)$ is represented by an element $x \in \mathcal{A}(L_{S,\epsilon_0})^E$ for some $E > 0$. Hence, by the previous lemma, there is a $C > 0$ such that

$$\alpha_i \circ \alpha_{i-1} \circ \ldots \circ \alpha_0(x) \in \mathcal{A}(L_{S,\epsilon_{i+1}})^{CE}$$

for all $i \geq 0$. It follows that y is represented by some element in $\mathcal{A}(L_{S,\epsilon_i})^{E_i}$ for big enough $i \gg 0$. $\square$

We now consider the sequence

$$\ldots \xrightarrow{\beta_{i-1}} \mathcal{A}(L,S)^{E_i} \xrightarrow{\beta_i} \mathcal{A}(L,S)^{E_{i+1}} \xrightarrow{\beta_{i+1}} \ldots,$$

where

$$\beta_i := \Psi_{i+1} \circ \alpha_i \circ \Psi_i^{-1} \colon (\mathcal{A}(L,S)^{E_i}, \partial_{S,N}) \to (\mathcal{A}(L,S)^{E_{i+1}}, \partial_{S,N}),$$

and define

$$(\mathcal{A}(L,S),\partial_{S,N}) := \lim_{i\to\infty}(\mathcal{A}(L,S)^{E_i},\partial_{S,N})$$

as the direct limit of this sequence.

Remark 6.3. One should expect that this limit is the same as the limit with respect to sequence of inclusion maps

$$\ldots \subset (\mathcal{A}(L,S)^{E_i},\partial_{S,N}) \subset (\mathcal{A}(L,S)^{E_{i+1}},\partial_{S,N}) \subset \ldots,$$

since the maps in the definition of β_i should stabilize below each fixed action level. However, we do not prove this.

The maps

$$\Phi_i\colon\ (\mathcal{A}(L_{S,\epsilon_i})^{E_i},\partial_i) \to (\mathcal{A}(L),\partial)$$

induced by the exact Lagrangian cobordism V_{ϵ_i} satisfy the equality

$$\Phi_i = \Phi_{i+1}\circ\alpha_i\colon\ H(\mathcal{A}(L_{S,\epsilon_i,\eta_i})^{E_i},\partial_i) \to H(\mathcal{A}(L),\partial)$$

on homology level (see Section 2.5). Hence, there is an induced map

$$\Phi\colon\ \lim_{i\to\infty} H(\mathcal{A}(L_{S,\epsilon_i})^{E_i},\partial_i) \to H(\mathcal{A}(L),\partial).$$

Theorem 6.4. *There is a DGA-isomorphism*

$$\Psi\colon\ \lim_{i\to\infty}(\mathcal{A}(L_{S,\epsilon_i})^{E_i},\partial_i) \to (\mathcal{A}(L,S),\partial_{S,N}).$$

Moreover, the map

$$\Phi\colon\ \lim_{i\to\infty} H(\mathcal{A}(L_{S,\epsilon_i})^{E_i},\partial_i) \to H(\mathcal{A}(L),\partial)$$

induced by the exact Lagrangian cobordism V_ϵ has the property that

$$\Phi\circ\Psi^{-1}\colon\ H(\mathcal{A}(L,S)^{E_i},\partial_{S,N}) \to H(\mathcal{A}(L),\partial)$$

is induced by the quotient projection

$$\lim_{i\to\infty}(\mathcal{A}(L,S)^{E_i},\partial_{S,N}) \to \lim_{i\to\infty}(\mathcal{A}(L,S)/\langle s\rangle^{E_i},\partial_{S,N}).$$

Proof. The first claim follows by studying the commutative diagram

$$\begin{tikzcd}
\ldots \arrow[r, "\alpha_{i-1}"] & \mathcal{A}(L_{S,\epsilon_i})^{E_i} \arrow[r, "\alpha_i"] \arrow[d, "\Psi_i", "\simeq"'] & \mathcal{A}(L_{S,\epsilon_{i+1}})^{E_{i+1}} \arrow[r, "\alpha_{i+1}"] \arrow[d, "\Psi_{i+1}", "\simeq"'] & \ldots \\
\ldots \arrow[r, "\beta_{i-1}"] & \mathcal{A}(L,S)^{E_i} \arrow[r, "\beta_i"] & \mathcal{A}(L,S)^{E_{i+1}} \arrow[r, "\beta_{i+1}"] & \ldots,
\end{tikzcd}$$

where $\beta_i = \Psi_{i+1}\circ\alpha_i\circ\Psi_i^{-1}$, and where all Ψ_i are isomorphisms of DGAs.

For the other claims, recall that

$$\Phi_i\circ\Psi_i^{-1}\colon\ (\mathcal{A}(L,S)^{E_i},\partial_{S,N}) \to (\mathcal{A}(L)^{E_i},\partial)$$

corresponds to the quotient projection

$$(\mathcal{A}(L,S)^{E_i},\partial_{S,N}) \to (\mathcal{A}(L,S)^{E_i}/\langle s\rangle,\partial_{S,N}),$$

and that the equality

$$\begin{aligned} &\Phi_i \circ \Psi_i^{-1} = \\ &= \quad (\Phi_{i+1} \circ \alpha_i) \circ \Psi_i^{-1} = (\Phi_{i+1} \circ \Psi_{i+1}^{-1}) \circ (\Psi_{i+1} \circ \alpha_i \circ \Psi_i^{-1}) \\ &= \quad (\Phi_{i+1} \circ \Psi_{i+1}^{-1}) \circ \beta_i \end{aligned}$$

holds on homology level. □

Corollary 6.5. *If $(\mathcal{A}(L_S), \partial)$ has an augmentation, then so does the DGA $\lim_{i\to\infty} \left(\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i\right)$ and, in particular, so does $(\mathcal{A}(L,S), \partial_{S,N})$. Moreover, it follows that $(\mathcal{A}(L), \partial)$ has an augmentation as well.*

Proof. Suppose that $(\mathcal{A}(L_S), \partial)$ has an augmentation. It follows that the DGA

$$\left(\lim_{i\to\infty} (\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i\right)$$

has an augmentation as well, since the set

$$Aug\left(\lim_{i\to\infty} (\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i)\right)$$

of augmentations is the inverse limit of the sequence

$$\ldots \xleftarrow{\alpha_{i-1}^*} Aug((\mathcal{A}(L_{S,\epsilon_i})^{E_i}, \partial_i)) \xleftarrow{\alpha_i^*} Aug((\mathcal{A}(L_{S,\epsilon_{i+1}})^{E_{i+1}}, \partial_{i+1})) \xleftarrow{\alpha_{i+1}^*} \ldots,$$

and since all of the sets in the sequence are finite and non-empty (that is, it satisfies a kind of Mittag-Leffler condition).

The previous theorem immediately implies that $(\mathcal{A}(L,S), \partial_{S,N})$ has an augmentation. In particular, for each i, the DGA $(\mathcal{A}(L,S)^{E_i}, \partial_{S,N,i})$ has an augmentation, and hence, by Corollary 1.5, so does each

$$(\mathcal{A}(L)^{E_i}, \partial_i) \simeq (\mathcal{A}(L,S)^{E_i}/\langle s\rangle, \partial_{S,N,i}).$$

It follows that there is a sequence

$$\ldots \leftarrow Aug((\mathcal{A}(L)^{E_i}, \partial_i)) \leftarrow Aug((\mathcal{A}(L)^{E_{i+1}}, \partial_{i+1})) \leftarrow \ldots,$$

in which all sets are finite and non-empty. Again, this implies that

$$Aug(\mathcal{A}(L), \partial) = Aug\left(\lim_{i\to\infty} (\mathcal{A}(L)^{E_i}, \partial)\right) \neq \emptyset.$$

□

7. The case $k = n - 1$

For $(n-1)$-surgeries on n-dimensional Legendrian submanifolds the situation is completely different. Since every pseudo-holomorphic disk which has a boundary point mapping to $\mathbb{R} \times S$ is tangent to $\exp_{\mathbb{R}\times S}(\mathbb{R}\mathbf{v})$ of any given order at that point, it does not make sense to count these tangencies. We will modify the algebra to obtain results similar to Theorem 1.2 and 1.4, but where we impose some additional algebraic relations by hand. In a forthcoming work we will show that the results in this section are sufficient for providing formulas for the *Linearized Legendrian contact homology* of L_S in terms of data on L.

We assume that $S \subset L$ is a framed, and thus co-oriented, $(n-1)$-sphere, and consider the algebra

$$\mathcal{A}(L, S) := \mathbb{Z}_2\langle \mathcal{Q}(L) \cup \{s, t\}\rangle / \langle st + 1, ts + 1\rangle,$$

where

$$|s| = |t| = 0.$$

After perturbing S we may assume that all rigid (up to translation) pseudo-holomorphic disks in $\mathbb{R} \times Y$ having boundary on $\mathbb{R} \times L$ intersect $\mathbb{R} \times S$ transversely. We also assume that there are no Reeb chords on S.

We can now define a differential ∂_S on $\mathcal{A}(L, S)$ by counting the same rigid disks as we do in the definition of the standard Legendrian contact homology DGA for L, but while keeping track of the boundary points intersecting S. More precisely, for any Reeb chord $a \in \mathcal{Q}(L)$ and any word of Reeb chords $\mathbf{b} = b_1 \cdot \ldots \cdot b_m$ on L, we decompose the moduli spaces $\mathcal{M}_{a;\mathbf{b};A}(L)$ satisfying $\dim \mathcal{M}_{a;\mathbf{b};A}(L) = 1$ into

$$\mathcal{M}_{a;\mathbf{b};A}(L) = \bigcup_{\mathbf{w} \in \mathbb{Z}^{m+1}} \widetilde{\mathcal{M}}_{a;\mathbf{b},\mathbf{w};A}(L),$$

where $\widetilde{\mathcal{M}}_{a;\mathbf{b},\mathbf{w},A}(L)$ consists of the subset of the disks in $\mathcal{M}_{a;\mathbf{b};A}(L)$ whose boundary arc between the i:th and $(i+1)$:th puncture, starting the count at the positive puncture, has intersection-number w_i with $\mathbb{R} \times S$. We now define

$$\begin{aligned} &\partial_S(s) = 0, \\ &\partial_S(a) := \sum_{\substack{\dim \mathcal{M}_{a,\mathbf{b};A}(L)=1 \\ \mathbf{w} \in \mathbb{Z}^{m+1}}} |\widetilde{\mathcal{M}}_{a;\mathbf{b},\mathbf{w};A}(L)/\mathbb{R}| s^{w_1} b_1 s^{w_2} \cdot \ldots \cdot s^{w_m} b_m s^{w_{m+1}}, \end{aligned}$$

and extend the definition to all of $\mathcal{A}(L, S)$ using the Leibniz rule.

Proposition 7.1. $(\mathcal{A}(L, S), \partial_S)$ *is a well-defined DGA, whose homotopy type is invariant under Legendrian isotopy.*

Proof. This follows by the standard proofs, after making the following observation. In 1-dimensional families we generically see cancellations of incidence conditions on S. However, since we are counting the algebraic number of incidence conditions, canceling pairs of points do not contribute to this count. □

We now assume that there are ambient surgery data for $S \subset L$ such that, moreover, there are no Reeb chords on $L \cup D_S$ starting or ending on the bounding disk D_S.

Remark 7.2. Observe that since D_S is Lagrangian, generically there are Reeb chords on $L \cup D_S$ starting or ending on D_S. We can, however, make sure that the assumptions are satisfied after a Legendrian isotopy of L.

First, we may assume that there are no Reeb chords starting or ending in a neighborhood of the origin of D_S. Thus, after a Legendrian isotopy of L which radially contracts S along D_S, and which has support in a small enough neighborhood of D_S, we may assume that the corresponding radially contracted bounding disk has no Reeb chords. A Legendrian surgery on this small bounding disk again yields L_S. Of course, the former isotopy introduces new Reeb chords on L.

By the construction in Section 3.4 we get an exact Lagrangian cobordism V_ϵ from L to L_S, where L_S is the Legendrian submanifold obtained from L after ambient surgery along S. Let c_{new} denote the new Reeb chord on L_S, and let $\widetilde{S} \subset V_\epsilon$ denote the core disk of the handle attachment V_ϵ, as defined in Section 5.1. Recall that $\widetilde{S}$ coincides with $(-\infty, -1) \times S$ outside of a compact subset.

After changing $\widetilde{S}$ via a compactly supported perturbation, we may assume that every rigid pseudo-holomorphic disk in $\mathbb{R} \times Y$ with boundary on V_ϵ intersects $\widetilde{S}$ transversely.

Let $A \in H_1(V_\epsilon)$, $a \in \mathcal{Q}(L_S)$ and $b_1 \cdot \ldots \cdot b_m$ be a word of Reeb chords on L. We decompose the moduli spaces $\mathcal{M}_{a;\mathbf{b};A}(V_\epsilon)$ satisfying $\dim \mathcal{M}_{a;\mathbf{b};A}(V_\epsilon) = 0$ into

$$\mathcal{M}_{a;\mathbf{b};A}(V_\epsilon) = \bigcup_{\mathbf{w} \in \mathbb{Z}^{m+1}} \widetilde{\mathcal{M}}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon),$$

where $\widetilde{\mathcal{M}}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon)$ consists of the subset of the disks in $\mathcal{M}_{a,\mathbf{b};A}(V_\epsilon)$ whose boundary arc between the i:th and $(i+1)$:th puncture, starting the count at the positive puncture, has intersection-number w_i with $\widetilde{S}$.

We now define a DGA morphism

$$\Psi \colon (\mathcal{A}(L_S), \partial) \to (\mathcal{A}(L,S), \partial_S)$$

defined on the generators by

$$\Psi(a) = \sum_{\substack{\dim \mathcal{M}_{a,\mathbf{b};A}(V_\epsilon)=0 \\ \mathbf{w} \in \mathbb{Z}^{m+1}}} |\widetilde{\mathcal{M}}_{a;\mathbf{b},\mathbf{w};A}(V_\epsilon)| s^{w_1} b_1 s^{w_2} \cdot \ldots \cdot s^{w_m} b_m s^{w_{m+1}},$$

and then extended to an algebra map.

Proposition 7.3. *The map*

$$\Psi \colon (\mathcal{A}(L_S), \partial) \to (\mathcal{A}(L,S), \partial_S)$$

is a well defined injective DGA morphism. Furthermore, the DGA morphism

$$\Phi \colon (\mathcal{A}(L_S), \partial) \to (\mathcal{A}(L), \partial)$$

induced by the exact Lagrangian cobordism V_ϵ is surjective with kernel being the two-sided ideal $\langle c_{new} - 1 \rangle$. Finally, Φ is obtained by the composition of Ψ with the quotient projection onto

$$(\mathcal{A}(L,S), \partial_S)/\langle s-1 \rangle \simeq (\mathcal{A}(L), \partial).$$

Proof. The chain map property follows by a standard argument.

To see the injectivity we proceed as follows. Consider the lift of Ψ to a map

$$\widetilde{\Psi}\colon \mathcal{A}(L_S) \to \mathbb{Z}_2\langle \mathcal{Q}(L) \cup \{s,t\}\rangle,$$

and an extension of $\widetilde{\Psi}$ to $\widetilde{\widetilde{\Psi}}$ to

$$\widetilde{\widetilde{\Psi}}\colon \mathcal{A}(L_S) * \langle \widetilde{t}\rangle \to \mathbb{Z}_2\langle \mathcal{Q}(L) \cup \{s,t\}\rangle,$$

by setting $\widetilde{\widetilde{\Psi}}(\widetilde{t}) = t$.

The disk counts in Lemma 5.5 and 5.7 shows that $\widetilde{\widetilde{\Psi}}$ is a tame isomorphism and that the pre-image of the ideal $\langle st-1, ts-1\rangle$ is $\langle c_{new}\widetilde{t}-1, \widetilde{t}c_{new}-1\rangle$. This implies that Ψ is injective.

The same disk-count also shows that Φ is surjective with kernel given by $\langle c_{new}-1\rangle$. □

Remark 7.4. There are circumstances when the above map Ψ will be an isomorphism onto the free subalgebra

$$\widetilde{\mathcal{A}}(L,S) := \mathbb{Z}_2\langle \mathcal{Q}(L) \cup \{s\}\rangle \subset \mathcal{A}(L,S).$$

For instance, under certain geometric conditions on the bounding disk D_S, one may conclude that holomorphic disks with boundary on L have boundary always intersecting S positively.

8. The moduli space of pseudo-holomorphic disks with jet-constraints on the boundary

Here we discuss the set-up of the moduli spaces, and sketch the transversality proofs. We use constructions from [EES3], [Dra] and [ES], and refer to those papers for functional-analytical details.

In the following we suppose that we have an $(n+1)$-dimensional exact Lagrangian cobordism $V \subset \mathbb{R} \times Y$ which coincides with the cylinders

$$((-\infty,-N) \times L_1) \cup ((N,+\infty) \times L_2)$$

outside of a compact set for some $N \gg 0$ and where thus $L_1, L_2 \subset Y$ are Legendrian submanifolds. We assume that the Legendrian submanifolds L_1 and L_2 are closed and chord generic.

Fix a $(k+1)$-dimensional submanifold $S \subset V$ satisfying $k+1 < n$, together with a non-vanishing vector-field $\mathbf{v}$ along S which is normal to S inside V. We require both S and $\mathbf{v}$ to be $\mathbb{R}$-invariant outside of a compact subset. Moreover, we require S to be disjoint from $\mathbb{R} \times c$ outside some compact subset, where c is a Reeb chord on L_2.

8.0.1. *A model for a conformal structure on the disk with boundary punctures.* Let $\dot{D} := D \setminus \{p_0, \ldots, p_m\}$ denote the unit disk with $m+1$ fixed boundary points removed. We call the puncture p_0 *special*, and we suppose that $p_1 < \ldots < p_m$ with respect to the ordering of $\partial D \setminus \{p_0\}$ induced by the orientation.

Let C_{m+1} denote the space of conformal structures on $\dot{D}$, where two conformal structures are identified if they differ by an orientation-preserving diffeomorphism fixing the special puncture. This space is diffeomorphic to $\mathbb{R}^{m-2}$ for $m \geq 2$ and to a point for $m \leq 2$.

In the case $m > 0$, for every conformal structure c, we now construct a disk $\Delta_{c,m+1} \subset \mathbb{C}$ whose conformal structure induced by the complex structure is equivalent to c.

Start with the set

$$\mathbb{R} \times [0, 2m] \subset \mathbb{C}$$

and remove $m-1$ slits of height 1 whose s-coordinates range from $s_i \in \mathbb{R}$ to $+\infty$, where $i = 1, \ldots, m-1$, as shown in Figure 3. In this way, for some $N \gg 0$, we obtain m strips on the right in $\Delta_{c,m+1}$ which coincide with $(N, +\infty) \times [i-1, i] \subset \mathbb{C}$ outside of a compact set, and one strip on the left which coincides with $(-\infty, -N) \times [0, 2m]$ outside of a compact set.

For every conformal structure c there are choices of $s_i \in \mathbb{R}$, $i = 1, \ldots, m-1$ for which the conformal structure on the disk $\Delta_{c,m+1} \subset \mathbb{C}$ induced by the complex structure is equivalent to c. Moreover, up to a translation $s_i \mapsto s_i + C$ for all i, the choices of s_i are unique with that property.

For $m > 0$ we fix a representative by a disk $\Delta_{c,m+1}$ for each conformal structure c, such that the choice depends smoothly on c. We also fix an orientation-preserving conformal diffeomorphism

$$f_c \colon \dot{D} \to \Delta_{c,m+1} \subset \mathbb{C}$$

which maps the special puncture to the end at $s = -\infty$. Furthermore, we require that the map f_c depends smoothly on c.



Figure 3. The model $\Delta_{c,m+1}$ of a conformal structure on the disk with $m+1$ punctures. The strip on the left has height $2m$, while the m strips on the right all have height 1.

8.1. **The configuration space.** In the following we fix capping paths for each Reeb-chord on both the positive and the negative Legendrian end of V. We also fix a cylindrical almost complex structure J. Let U_S be a neighborhood of $S \subset V$ which is cylindrical outside of a compact set and fix an embedding ι of a neighborhood of $U_S \subset \mathbb{R} \times Y$ into $\mathbb{C}^N$. (We will later make additional assumptions on this embedding.)

Let a be a Reeb chord on the Legendrian submanifold at the positive end of V and $\mathbf{b} = b_1 \cdot \ldots \cdot b_m$ a word of Reeb-chords on the Legendrian submanifold at the negative end of V. Let

$$\begin{cases} \mathbf{w} = (w_1, \ldots, w_{m+1}) \in (\mathbb{Z}_{\geq 0})^{m+1}, \\ d_1, \ldots, d_{m+1} \in \mathbb{Z}_{\geq 0}, \\ \mathbf{l} = ((l_1^1, \ldots, l_{d_1}^1), \ldots, (l_1^{m+1}, \ldots, l_{d_{m+1}}^{m+1})) \in (\mathbb{Z}_{\geq 0})^{d_1} \times \ldots \times (\mathbb{Z}_{\geq 0})^{d_{m+1}}, \end{cases}$$

be tuples satisfying

$$(l_1^i + 1) + \ldots + (l_{d_i}^i + 1) = w_i.$$

In addition, we define

$$d := d_1 + \ldots + d_{m+1}.$$

Finally, we let $A \in H_1(V; \mathbb{Z})$, and fix $\eta > 0$.

Recall that $\partial \dot{D}$ has a natural ordering as a subset of $\partial D \setminus \{p_0\}$. Let

$$(\partial \dot{D})^d_{\mathbf{l}} \subset (\partial \dot{D})^{d_1} \times \ldots \times (\partial \dot{D})^{d_{m+1}}$$

be the open and connected set consisting of tuples

$$((\theta_1^1, \ldots, \theta_{d_1}), \ldots, (\theta_1^{m+1}, \ldots, \theta_{d_{m+1}}^{m+1}))$$

satisfying the property that θ_j^i are increasing with respect to the lexicographic order on (i, j) and, moreover, satisfies $p_i < \theta_j^i < p_{i+1}$. Here we set $p_{m+1} := p_0$.

In other words, $(\partial \dot{D})^d_{\mathbf{l}}$ consists of a d-tuple of points in $\partial \dot{D}$ which come in increasing order (relative the order on $\partial D \setminus \{p_0\}$ induced by the orientation) and such that there are precisely d_i points on the oriented arc of $\partial \dot{D}$ having starting point p_i.

We are interested in the configuration space

$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V, S; J) \subset W^{2,2}_{loc}((\dot{D}, \partial \dot{D}), (\mathbb{R} \times Y, V)) \times C_{m+1} \times (\partial \dot{D})^d_{\mathbf{l}} \times (U_S)^d,$$

consisting of tuples $(g, c, (\theta_j^i), (s_j^i))$ satisfying the following conditions.

The boundary condition. The map $g \colon (\dot{D}, \partial \dot{D}) \to (\mathbb{R} \times Y, V)$ takes the boundary into V and is locally in the Sobolev space $W^{2,2}$ consisting of maps which are twice weakly differentiable and having differentials in L^2. This implies that g is continuous up to the boundary.

The point constraints. We require that $g(\theta_j^i) = s_j^i$.

The asymptotical properties. The conformal structure c on $\dot{D}$ induces a conformal structure c' on the disk

$$\ddot{D} := \dot{D} \setminus \cup \{\theta_j^i\}$$

having $m+1+d$ boundary punctures. In the case $m+d>0$, we identify this disk with the standard model $\Delta_{c',m+1+d}$ using $f_{c'}$ as defined above. Observe that this model has $m+1$ strips corresponding to punctures from c, and d strips corresponding to punctures at θ^i_j. We let (s,t) be coordinates on $\Delta_{c',m+1+d}$, as shown in Figure 3.

The asymptotics at the punctures p_i. By Darboux's theorem we may take contact-form preserving charts in Y around each of the Reeb chords $a, b_1, \ldots, b_m \subset Y$ on the form $((T^*\mathbb{R}^n)\times\mathbb{R}, dz-\sum_i p_i dq_i)$. By the assumption of chord genericity, the two branches of L_i at the endpoints of a Reeb chord intersect transversely when projected to $T^*\mathbb{R}^n$.

For a Reeb chord $b \subset Y$, we let

$$\rho_b\colon\ [0,\ell(c)] \to b$$

be its parametrization defined by $\dot{\rho_b} = R_{\rho_b}$. We require that there are constants $C_0,\ldots,C_m \in \mathbb{R}$ such that, relative the above coordinate charts and translations of the symplectization coordinate determined by the C_i, the map g has the following asymptotic behavior. Along the strip corresponding to the special puncture, we require the map $g=(g_\mathbb{R}, g_Y)$ to satisfy

$$e^{-\frac{\eta}{2m}s}\left(g_\mathbb{R}(s,t)-\frac{\ell(a)s}{2m}+C_0, g_Y(s,t)-\rho_a\left(\frac{\ell(a)(1-t)}{2m}\right)\right)\in W^{2,2},$$

and along the strip corresponding to the i:th non-special puncture, we require the map $g=(g_\mathbb{R}, g_Y)$ to satisfy

$$e^{\eta s}\left(g_\mathbb{R}(s,t)-\ell(b_i)s+C_i, g_Y(s,t)-\rho_{b_i}\left(\ell(b_i)(1-t)\right)\right)\in W^{2,2}.$$

The asymptotics at the points $\{\theta^i_j\}$. Along the strip of $\Delta_{c',m+1+d}$ corresponding to the boundary point θ^i_j, we require that the map g satisfies

$$e^{(\pi l^i_j+\eta)s}(\iota(g(s,t))-\iota(s^i_j))\in W^{2,2}.$$

The homology class. Consider the compactification $\overline{V}$ of V obtained by adding a copy of its Legendrian positive and negative end at $\pm\infty$, respectively. The map $g|_{\partial\dot{D}}$ extends to a union of continuous paths on $\overline{V}$. Concatenating these paths with the appropriate capping paths of the Reeb chords on the respective Legendrian ends closes up the boundary of the disk. We require the obtained cycle to be in homology class $A \in H_1(V;\mathbb{Z}) \simeq H_1(\overline{V},\mathbb{Z})$.

J-holomorphicity along the boundary. Let

$$\overline{\partial}_J g = dg + Jdg\circ i \in \mathcal{W}^{2,1,\eta}(T^{*0,1}\dot{D}\otimes g^*T(\mathbb{R}\times Y))$$

denote the Cauchy-Riemann operator with respect to the almost complex structure J on $\mathbb{R}\times Y$ and the complex structure i on $\dot{D}$ determined by c. The map g is then required to be J-holomorphic along the boundary in the following distributional sense:

$$\int_{\partial\dot{D}}\langle\overline{\partial}_J g, f\rangle = 0,$$

for all continuous sections f of the bundle

$$T^{*0,1}(D)\otimes g^*(T(\mathbb{R}\times Y))|_{\partial\dot{D}}\to\partial\dot{D}.$$

The space $\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$ has the structure of a Banach manifold. For more details, we refer to [EES3].

8.2. **The fiber-bundle structure of the configuration space.**

The natural projection

$$\pi\colon\ \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)\to(\partial\dot{D})^d_{\mathbf{l}}\times(U_S)^d$$

gives the Banach manifold $\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$ the structure of the total-space of a fiber-bundle. To construct local trivializations, we proceed as in [ES, §4.4].

The problem that one has to take care of is to translate the disks in a given fibre to disks in the nearby fibers, while keeping the holomorphicity along the boundary. Consider the fiber over $(\theta^i_j,s^i_j)\in(\partial\dot{D})^d_{\mathbf{l}}\times(U_S)^d$. For (θ'^i_j,s'^i_j) close enough to (θ^i_j,s^i_j) we have to construct a compactly supported diffeomorphism

$$\psi_{(s'^i_j)}\colon\ \mathbb{R}\times Y\to\mathbb{R}\times Y$$

depending smoothly on (s'^i_j) and satisfying

(1) $\psi_{(s'^i_j)}(s^i_j)=s'^i_j$.
(2) $\psi_{(s'^i_j)}(V)=V$.
(3) $\psi_{(s'^i_j)}$ is J-holomorphic along V.

For the construction of $\psi_{(\theta'^i_j,s'^i_j)}$ we refer to [ES] where such a map is constructing by using linear translations and cut-off functions which are J-holomorphic along V.

We also have to construct diffeomorphisms

$$\varphi_{(\theta'^i_j)}\colon\ \dot{D}\to\dot{D},$$

depending smoothly on θ'^i_j and satisfying

(1) $\varphi_{(\theta'^i_j)}(\theta^i_j)=\theta'^i_j$.
(2) $\varphi_{(\theta'^i_j)}(\partial\dot{D})=\partial\dot{D}$.
(3) $\varphi_{(\theta'^i_j)}$ is holomorphic along $\partial\dot{D}$.
(4) $\varphi_{(\theta'^i_j)}$ is holomorphic in a neighborhood of $\{\theta^i_j\}\subset\partial\dot{D}$.
(5) $\varphi_{(\theta'^i_j)}=id_{\dot{D}}$ outside of a compact set.

We can now translate the fiber over (θ^i_j,s^i_j) to the fiber over (θ'^i_j,s'^i_j) by simply mapping $g\mapsto\psi_{(s'^i_j)}\circ g\circ\varphi_{(\theta'^i_j)}$.

8.3. **The $\overline{\partial}$-operator.** We let

$$\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J),$$

denote the fiber bundle whose fiber above $(g,c,(\theta^i_j),(s^i_j))$ consists of sections Γ of the bundle

$$T^{*0,1}(D)\otimes g^*(T(\mathbb{R}\times Y)) \to \dot{D},$$

such that Γ is in the Sobolev space $W^{1,2,\eta}$ with one derivative and positive weights at punctures $p_0,\ldots,p_m$ defined analogously as for the spaces $\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$.

Moreover, we require that Γ satisfies the following vanishing conditions at the points θ^i_j. The embedding ι of a neighborhood of $U_S\subset\mathbb{R}\times Y$ into $\mathbb{C}^N$ induces a push-forward map

$$\iota_*\colon\ T^{*0,1}\dot{D}\otimes g^*(T(R\times Y)) \to T^{*0,1}\dot{D}\otimes\mathbb{C}^N$$

defined at the points in $\dot{D}$ which are mapped into this neighborhood of U_S by g. We require Γ to satisfy

$$e^{(\pi l^i_j+\eta)s}(\iota_*(\Gamma(s,t)))\in W^{2,1}$$

along the infinite strip in $\Delta_{c,m+1+d}$ corresponding to the point $\theta^i_j\in\partial\dot{D}$.

Finally, we require that Γ vanishes along the boundary in sense that

$$\int_{\partial\dot{D}}\langle\Gamma,f\rangle=0,$$

for all continuous sections f of the bundle

$$T^{*0,1}(D)\otimes g^*(T(\mathbb{R}\times Y))|_{\partial\dot{D}}\to\partial\dot{D}.$$

The operator

$$\overline{\partial}(g)=dg+J\circ dg\circ i$$

acts on functions from a Riemann-surface into $\mathbb{R}\times Y$. In our setting it extends to a section

$$\overline{\partial}\colon\ \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)\to\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J).$$

That this operator is Fredholm for $\eta>0$ small enough follows from the results in [EES3].

We define the moduli space

$$\mathcal{M}_{a;\mathbf{b};A}(V):=\overline{\partial}^{-1}(0)\subset\mathcal{H}^{2,\eta}_{a;\mathbf{b},0;0;A}(V,S;J).$$

8.4. **A holomorphic neighborhood of S.** For the following constructions we have to make additional assumptions on the almost complex structure. We will assume that J is an almost complex structure on the symplectization $\mathbb{R}\times Y$ which is compatible with the symplectic form, cylindrical outside of a compact set, and which satisfies the following conditions. Every point of S should have a neighborhood in $\mathbb{R}\times Y$ which is biholomorphic to a neighborhood of the origin of

$$\mathbb{C}^{n+1}\simeq T^*(\mathbb{R}^k\times\mathbb{R}\times\mathbb{R}^{n-k-1})$$

endowed with the standard complex structure such that, moreover, the following real-analyticity conditions for V, S and $\mathbf{v}$ are fulfilled:

- The zero-section is identified with V.
- The subset $\mathbb{R}^{k+1} \times \{0\}$ of the zero-section is identified with S.
- The vector field $\mathbf{v}$ is identified with the coordinate vector-field corresponding to the $(k+2)$:th $\mathbb{R}$-factor of the zero-section.

Observe that in the case when, $V = \mathbb{R} \times L$, $S = \mathbb{R} \times T$, and $\mathbf{v}$ all are $\mathbb{R}$-invariant, there always is such a cylindrical almost complex structure on $\mathbb{R} \times Y$. In the non-cylindrical case, one may have to drop the requirement of $\mathbb{R}$-invariance to guarantee the existence.

We fix a neighborhood U_S of $S \subset V$ which can be covered by charts of the above form. Using the Whitney embedding theorem for analytic manifolds, we can find an analytic embedding $\iota\colon U_S \hookrightarrow \mathbb{R}^N$. Moreover, there is an analytic retract $r\colon N(U_S) \to U_S$ from a tubular neighborhood of $\iota(U_S) \subset \mathbb{R}^N$.

Observe that the above analytic embedding $\iota\colon U_S \hookrightarrow \mathbb{R}^N$ extends to a holomorphic embedding of a neighborhood of $U_S \subset \mathbb{R} \times Y$ into $\mathbb{C}^N$ and that the retract r is extended to a holomorphic retract onto the image of this embedding. By abuse of notation, we will use ι and r to denote the extensions of the embedding and retract, respectively, as well.

In this case, we will assume that the embedding ι in the construction of $\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$ coincides with the above holomorphic embedding. Moreover, we will assume that the map $\psi_{(s'^i_j)}$ constructed in Section 8.2 is holomorphic in a neighborhood of $\{s^i_j\} \subset \mathbb{R} \times Y$.

8.5. **The configuration space with jet conditions.** Let $J^l(\partial D, X)$ be the space of l-jets of maps $\partial D \to X$ into a smooth manifold X. This is a smooth fiber-bundle with base $\partial D \times X$ whose fiber above (θ, x) consists of equivalence classes of germs of smooth maps $f\colon \partial D \to X$ satisfying $f(\theta) = x$, where two germs are considered to be equivalent if their Taylor expansions agree up to order l at θ. We have an identification

$$J^l(\partial D, X) \simeq \partial D \times (\underbrace{TX \oplus \ldots \oplus TX}_{l}) \to \partial D \times X.$$

Denote

$$J_i := J^{l^i_1}(\partial \dot{D}, U_S) \times \ldots \times J^{l^i_{d_i}}(\partial \dot{D}, U_S), \;\; i = 1, \ldots, m+1$$

and consider the the bundle

$$p\colon \; J^{\mathbf{l}} := J_1 \times \ldots \times J_{m+1}|_{(\partial \dot{D})^d_{\mathbf{l}} \times (U_S)^d} \to (\partial \dot{D})^d_{\mathbf{l}} \times (U_S)^d.$$

obtained by restriction.

The configuration space with jet conditions consists of the pullback bundle $\pi^* J^{\mathbf{l}}$, where we recall that

$$\pi\colon \; \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to (\partial \dot{D})^d_{\mathbf{l}} \times U_S^d$$

is a fiber bundle in itself. This space will be denoted by

$$\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) := \pi^* J^{\mathbf{l}} \to \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J).$$

Observe that $\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$ is naturally a bundle over $(\partial\dot{D})^d_{\mathbf{l}} \times (U_S)^d$ as well.

A point in $\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$ is described by a tuple $(g,c,(\theta^i_j),(s^i_j),(g^i_j))$ where c is a conformal structure on $\dot{D}$, g is a map $\dot{D} \to \mathbb{R}\times Y$, $\theta^i_j \in \partial\dot{D}$, $g(\theta^i_j) = s^i_j \in U_S$, and g^i_j are l^i_j-jets of germs of smooth functions $\partial\dot{D} \to U_S$ mapping θ^i_j to s^i_j.

8.6. **Parameterizing maps by jet conditions.**

Define

$$\begin{aligned}
\widetilde{\mathbf{w}} &:= (d_1,\ldots,d_{m+1}) \in (\mathbb{Z}_{\geq 0})^{m+1},\\
\widetilde{\mathbf{l}} &:= ((0,\ldots,0),\ldots,(0,\ldots,0)) \in \mathbb{Z}^{d_1}_{\geq 0} \times \ldots \times \mathbb{Z}^{d_{m+1}}_{\geq 0},
\end{aligned}$$

and observe that there is an inclusion

$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \subset \mathcal{H}^{2,\eta}_{a;\mathbf{b},\widetilde{\mathbf{w}};\widetilde{\mathbf{l}};A}(V,S;J).$$

Lemma 8.1. *There is a smooth map*

$$\begin{aligned}
\Psi\colon\ &\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{H}^{2,\eta}_{a;\mathbf{b},\widetilde{\mathbf{w}};\widetilde{\mathbf{l}};A}(V,S;J),\\
&(g,c,(\theta^i_j),(s^i_j),(g^i_j)) \mapsto (\widetilde{g},c,(\theta^i_j),(s^i_j)),
\end{aligned}$$

of bundles over $(\partial\dot{D})^d_{\mathbf{l}} \times (U_S)^d$ which has the property that

1. *The restriction of Ψ to the zero-section*
$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \subset \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$$
is the inclusion map $\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \subset \mathcal{H}^{2,\eta}_{a;\mathbf{b},\widetilde{\mathbf{w}};\widetilde{\mathbf{l}};A}(V,S;J)$.
2. *The l^i_j-jet of $\widetilde{g}$ at θ^i_j is equal to g^i_j.*
3. *Let ι be the holomorphic embedding of a neighborhood of U_S into $\mathbb{C}^N$. The difference $\overline{\partial}(\iota\circ\widetilde{g} - \iota\circ g)$ has compact support in $\dot{D}\setminus\{\theta^i_j\}$.*
4. *The differential $D\Psi(\partial_{g^i_j})$, where $\partial_{g^i_j}$ is a variation of the germ g^i_j, is a section over $\dot{D}$ which is J-holomorphic outside of a relatively compact set in $\dot{D}\setminus\{\theta^i_j\}$.*

Proof. We fix a point $(g,c,(\theta^i_j),(s^i_j),(g^i_j))$. To define

$$\Psi(g,c,(\theta^i_j),(s^i_j),(g^i_j)) = (\widetilde{g},c,(\theta^i_j),(s^i_j)),$$

it suffices to specify the map $\widetilde{g}\colon \dot{D} \to \mathbb{R}\times Y$.

By assumption, we can choose a locally finite open cover $\{U_k\}$ of $U_S \subset V$ by Euclidean analytic neighborhoods obtained from intersections $U_i = \widetilde{U}_k \cap U_S$, where $\widetilde{U}_k$ is an Euclidean holomorphic neighborhood inside $\mathbb{R}\times Y$ for which U_S, S and $\mathbf{v}$ satisfy real-analyticity conditions. Also, we fix a partition of unity $\{\phi_k : \widetilde{U}_k \to \mathbb{R}\}$ subordinate to $\{\widetilde{U}_k\}$, which thus restricts to a partition of unity subordinate to $\{U_k\}$.

For each θ^i_j we do the following construction. We conformally identify $\dot{D}$ with the unit disk endowed with the standard complex structure which we, in turn, holomorphically identify with the upper half-plane $\{z = (x, y),\ y \geq 0\}$ via the map

$$D^2 \ni z \mapsto -i \frac{z - e^{i\theta^i_j}}{z + e^{i\theta^i_j}}.$$

The coordinate function x on the upper half-plane thus becomes an induced analytic coordinate on $\partial\dot{D}$ under which θ^i_j corresponds to 0.

Pick a representative of the l^i_j-jet g^i_j given by a function $g^i_{j,k} \colon \mathbb{R} \to U_S$ which is defined uniquely by the requirement that it is given by a polynomial of degree at most l^i_j in the analytic coordinate x relative the fixed analytic chart on U_k.

Recall that there is an analytic embedding ι of U_S into $\mathbb{R}^N$, together with a retract r onto $\iota(U_S)$. Now consider the real analytic function

$$\widetilde{g}^i_j(x) := r\left(\sum_k \phi_k(s^i_j)\left(\iota \circ g^i_{j,k}(x)\right)\right)$$

which is defined in a neighborhood of $\theta^i_j \in \partial\dot{D}$ and which has germ g^i_j. The function $\widetilde{g}^i_j$ depends smoothly on $(c, (\theta^i_j), (s^i_j), (g^i_j))$.

Observe that we can extend $\widetilde{g}^i_j$ to a unique holomorphic function defined on some neighborhood of the origin in the upper half-plane having image in $\mathbb{R} \times Y$.

Conformally identify $\dot{D} \backslash \cup \{\theta^i_j\}$ with $\Delta_{c',m+1+d}$ as in Section 8.1. Let (s, t) be coordinates on $\Delta_{c',m+1+d}$ as shown in Figure 3.

We let $a^i_j > 0$ be a real number depending smoothly on $(g, c, (\theta^i_j), (s^i_j), (g^i_j))$ such that the map $\widetilde{g}^i_j$, defined on the boundary of the strip in $\Delta_{c',m+1+d}$ corresponding to θ^i_j, extends to a holomorphic function defined for $s \geq a^i_j$ on that strip.

We fix smooth cut-off functions $\alpha^i_j(s, t)$ defined on the infinite strip in $\Delta_{c',m+1+d}$ corresponding to θ^i_j which satisfy

$$\alpha^i_j(s,t) = \begin{cases} 0, & s \leq 0 \\ 1, & s \gg 0, \end{cases}$$

and which are holomorphic on the boundary of the respective strip.

There is an real-analytic exponential map $\exp\colon TU_S \to U_S$, defined using a real-analytic metric, which extends to a holomorphic exponential map defined on some neighborhood of the zero-section of $\mathbb{C} \otimes TU_S$.

On the strip corresponding to θ^i_j, we now define $\widetilde{g}(s, t)$ by

$$\widetilde{g}(s,t) := \exp_{s^i_j}\left(\exp^{-1}_{s^i_j}(g(s,t)) + \alpha^i_j(s - a^i_j, t)(\exp^{-1}_{s^i_j}(\widetilde{g}^i_j(s,t))\right),$$

which is well-defined given that $a^i_j \gg 0$ is big enough. Finally, we extend $\widetilde{g}$ by g to the rest of $\Delta_{c',m+1+d}$. □

By composition we get a map

$$G := \overline{\partial} \circ \Psi \colon \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{E}^{1,\eta}_{a;\mathbf{b},\widetilde{\mathbf{w}};\widetilde{\mathbf{l}};A}(V,S;J).$$

Lemma 8.2. *The map G has image in*

$$\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w},\mathbf{l};A}(V,S;J) \subset \mathcal{E}^{1,\eta}_{a;\mathbf{b},\widetilde{\mathbf{w}};\widetilde{\mathbf{l}};A}(V,S;J).$$

Moreover,

$$G \colon \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w},\mathbf{l};A}(V,S;J)$$

has the property that the image of the linearized operator $D_{(g,c,(\theta^i_j),(s^i_j),(g^i_j))}G$ at the point $(g,c,(\theta^i_j),(s^i_j),(g^i_j))$ is a compact perturbation of the Fredholm operator $D_{(g,c,(\theta^i_j),(s^i_j))}\overline{\partial}$ at the point $(g,c,(\theta^i_j),(s^i_j))$, where

$$\overline{\partial} \colon \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J).$$

In particular, G is Fredholm as well.

Proof. Consider the point $(g,c,(\theta^i_j),(s^i_j),(g^i_j)) \in \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$, and let

$$\Psi(g,c,(\theta^i_j),(s^i_j),(g^i_j)) = (\widetilde{g},c,(\theta^i_j),(s^i_j)).$$

By part (3) of Lemma 8.1, it follows that the sections $\iota_*(\overline{\partial}\widetilde{g})$ and $\iota_*(\overline{\partial}g)$ in $T^{*0,1}\dot{D} \otimes \mathbb{C}^N$, which are defined in a neigbhourhood of the points $\{\theta^i_j\}$, satisfy

$$\iota_*(\overline{\partial}\widetilde{g}) = \iota_*(\overline{\partial}g)$$

in some neighborhood of θ^i_j. Since $\overline{\partial}g \in \mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w},\mathbf{l};A}(V,S;J)$, the same is true for $\overline{\partial}\widetilde{g}$.

The second statement follows by part (3) of Lemma 8.1 as well, since g and $\widetilde{g}$ coincide outside of some neighborhood of $\{\theta^i_j\}$, and since we may assume that

$$\iota_*(\overline{\partial}\widetilde{g}) = \iota_*(\overline{\partial}g) + \overline{\partial}\alpha$$

holds inside this neighborhood, where $\alpha(g,c,(\theta^i_j),(s^i_j),(g^i_j))$ is a smooth $\mathbb{C}^N$-valued function which is holomorphic outside of a compact subset of $\dot{D}$. □

Define

$$\mathcal{M}_{a;\mathbf{b};A}(V)_{U_S} \subset M_{a;\mathbf{b};A}(V) \times (\partial\dot{D})^d_{\mathbf{l}}$$

to be the open submanifold consisting of points $(g,(\theta^i_j))$ for which $g(\theta^i_j) \in U_S$.

Proposition 8.3. *Suppose that $\mathcal{M}_{a;\mathbf{b};A}(V)$ is transversely cut out. The map*

$$\Psi \colon \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{H}^{2,\eta}_{a;\mathbf{b},\widetilde{\mathbf{w}};\widetilde{\mathbf{l}};A}(V,S;J)$$

can be constructed so that the restriction

$$\begin{aligned}&\Psi|_{G^{-1}(0)}\colon\ G^{-1}(0)\to\mathcal{M}_{a;\mathbf{b};A}(V)_{U_S},\\&(g,c,(\theta^i_j),(g(\theta^i_j)),(g^i_j))\mapsto(\widetilde{g},\theta^i_j).\end{aligned}$$

is a diffeomorphism.

Proof. Consider the cut-off functions $\alpha^i_j(s-a^i_j,t)$ which are defined in the proof of Lemma 8.1. These are defined on the infinite strips of the disk

$$\Delta_{c',m+1+d}$$

corresponding to the points θ^i_j, and are used in the construction of Ψ. The translations a^i_j of the s-coordinates of these cut-off functions depend on the point in $\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$. The statement of the proposition will follow after choosing a^i_j with some care.

Fix a conformal structure c' on the $(m+1+d)$-punctured disk. Consider the set

$$C_{c'}\subset\mathcal{M}_{a;\mathbf{b};A}(V)\times(\partial\dot{D})^d_{\mathbf{l}}$$

consisting of points $(g,(\theta^i_j))$ for which the induced conformal structure on $\dot{D}\setminus\{\theta^i_j\}$ is equal to c'. By the compactness theorem in [BEH$^+$], this is a compact subset, since no bubbling can occur without changing the conformal structure.

The compactness of $C_{c'}$ implies that the map sending

$$(g,(\theta^i_j))\in C_{c'}\cap\mathcal{M}_{a;\mathbf{b};A}(V)_{U_S}$$

to the l^i_j-jet of g at θ^i_j has a precompact image $K_{c'}\subset J^{\mathbf{l}}$.

For each fixed conformal structure c_0 and c'_0, we may now choose a^i_j in the construction of Ψ to be constant on the set consisting of points

$$(g,c,(\theta^i_j),(g(\theta^i_j)),(g^i_j))\in\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$$

satisfying the following properties, where $\epsilon>0$ is some small number which is allowed to depend on c and c'_0.

- $c=c_0$
- The induced conformal structure on $\dot{D}\setminus\{\theta^i_j\}$ is c'_0.
- $(g,(\theta^i_j))$ is uniformly ϵ-close to $C_{c'}$.
- $(g^i_j)\in K_{c'}$.

We can moreover assume that $\widetilde{g}$ constructed using these functions a^i_j is uniformly ϵ-close to g.

With a^i_j satisfying the above properties the statement follows. We give the proof for injectivity. Suppose that

$$\Psi(g,c,(\theta^i_j),(g(\theta^i_j)),(g^i_j))=\Psi(g',d,(\theta'^i_j),(g(\theta'^i_j)),(g'^i_j))\in\mathcal{M}_{a;\mathbf{b};A}(V)_{U_S}.$$

Clearly, the points must satisfy

$$\begin{cases} c = d, \\ (\theta^i_j) = (\theta'^i_j), \\ s^i_j := g(\theta^i_j)) = (g(\theta'^i_j)), \\ (g^i_j) = (g'^i_j) \in K_{c'}, \\ \widetilde{g} = \widetilde{g'}. \end{cases}$$

Moreover, both $(g, (\theta^i_j))$ and $(g', (\theta'^i_j))$ are uniformly ϵ-close to $C_{c'}$. By construction, it now follows that

$$\exp^{-1}_{s^i_j}(\widetilde{g}) - \exp^{-1}_{s^i_j}(g) = \exp^{-1}_{s^i_j}(\widetilde{g'}) - \exp^{-1}_{s^i_j}(g'),$$

and hence that $g = g'$. Here exp is the holomorphic extension of the exponential map, as in the proof of Lemma 8.1. Surjectivity follows by a similar argument.

We leave out the proof that the restriction $\Psi|_{G^{-1}(0)}$ is a diffeomorphism. □

8.7. **The definition of the general moduli spaces.** Let

$$Z_{\mathbf{l},\mathbf{v}} \subset p^{-1}((\partial\dot{D})^d_{\mathbf{l}} \times S^d) \subset J^{\mathbf{l}}$$

be the manifold with corners defined by the property that $Z_{\mathbf{l},\mathbf{v}}$ consists of tuples of germs

$$((g^1_1, \ldots, g^1_{d_1}), \ldots, (g^{m+1}_1, \ldots, g^{m+1}_{d_{m+1}})),$$

where g^i_j is the l^i_j-jet of the germ of a map $\partial\dot{D} \to U_S$ which is tangent of order l^i_j to

$$\exp_S(\mathbb{R}\mathbf{v})$$

such that, in the case $l^i_j > 0$, the tangent of the germ has a non-negative component in the $\mathbf{v}$-direction. Here, exp is defined using some choice of metric on V.

We define $\widetilde{Z}_{\mathbf{l},\mathbf{v}}$ to be the boundary of $Z_{\mathbf{l},\mathbf{v}}$. This is a subvariety of $Z_{\mathbf{l},\mathbf{v}}$ with the additional constraint that there is least one index (i, j) for which $l^i_j > 0$ and whose corresponding germ g^i_j is tangent to S.

One can check that $Z_{\mathbf{l},\mathbf{v}}$ is a submanifold of codimension

$$(n-k)\sum_{i=1}^{m+1} d_i + (n-k-1)\sum_{i=1}^{m+1}\sum_{j=1}^{d_i} l^i_j,$$

with corners and that $\widetilde{Z}_{\mathbf{l},\mathbf{v}} \subset Z_{\mathbf{l},\mathbf{v}}$ is a subvariety of codimension 1.

Recall that we have the following pull-back diagram of vector-bundles.

$$\begin{tikzcd}
\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \arrow[r, "\widetilde{\pi}"] \arrow[d] & J^{\mathbf{l}} \arrow[d, "p"] \\
\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \arrow[r, "\pi"] & (\partial\dot{D})^d_{\mathbf{l}} \times (U_S)^d
\end{tikzcd}$$

In the following, we will denote the induced natural bundle map by

$$ev^{\mathbf{l}} := \widetilde{\pi} \colon \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to J^{\mathbf{l}}.$$

We start by defining the moduli spaces

$$\begin{aligned}
\mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v}) &:= (G, ev^{\mathbf{l}})^{-1}(\{0\} \times Z_{\mathbf{l},\mathbf{v}}),\\
\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v}) &:= (G, ev^{\mathbf{l}})^{-1}(\{0\} \times \widetilde{Z}_{\mathbf{l},\mathbf{v}}).
\end{aligned}$$

Observe that in the generic situation Proposition 8.3 implies that the solutions in $\mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$ are in bijection with J-holomorphic maps $\dot{D} \to \mathbb{R}\times Y$ having boundary on V in the specified homology class, punctures at the Reeb chords $a, b_1, \ldots, b_m$ and l^i_j-fold tangencies to $\exp_S(\mathbb{R}\mathbf{v})$ at points $\theta^i_j \in \partial\dot{D}$ mapping to S, such that the tangent of the germ g^i_j has non-negative component in the $\mathbf{v}$-direction in the case $l^i_j > 0$.

The space $\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$ consists of solutions as above, but with the additional constraint that the differential along the boundary vanishes in the direction $\mathbf{v}$ at some point θ^i_j mapping to S and satisfying $l^i_j > 0$.

Finally, we define the full moduli spaces

$$\begin{aligned}
\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) &:= \bigcup_{\mathbf{l}} \mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v}),\\
\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) &:= \bigcup_{\mathbf{l}} \mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}),
\end{aligned}$$

consisting of the union of the moduli spaces for all possible choices of tuples

$$\begin{aligned}
&d_1, \ldots, d_{m+1} \in \mathbb{Z}_{\geq 0},\\
&\mathbf{l} = ((l^1_1, \ldots, l^1_{d_1}), \ldots, (l^{m+1}_1, \ldots, l^{m+1}_{d_{m+1}})) \in (\mathbb{Z}_{\geq 0})^{d_1} \times \ldots \times (\mathbb{Z}_{\geq 0})^{d_{m+1}},
\end{aligned}$$

for which

$$(l^i_1 + 1) + \ldots + (l^i_{d_i} + 1) = w_i.$$

Again, Proposition 8.3 implies that there is an inclusion

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial\dot{D})^d_{\mathbf{l}}$$

in the case when $\mathcal{M}_{a;\mathbf{b};A}(V)$ is transversely cut out.

8.8. **The expected dimension of the moduli spaces.** In this section we will show that the dimension formulas

$$\dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) = |a| - |b_1| - \ldots - |b_m| - |s| \sum_{i=1}^{m+1} w_i + \mu(A),$$
$$\dim \mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) < |a| - |b_1| - \ldots - |b_m| - |s| \sum_{i=1}^{m+1} w_i + \mu(A),$$

hold in the case whenever the respective solution spaces are transversely cut out, and when the weight $\eta > 0$ is chosen small enough.

The linearized $\overline{\partial}$-operator restricted to a fiber of

$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to (\partial \dot{D})^d_{\mathbf{l}} \times (U_S)^d$$

is the Cauchy-Riemann operator on a weighted Sobolev space, with weights at the punctures $\{p_i\} \cup \{\theta^i_j\} \subset \partial D$ depending on $\eta > 0$ as specified in Section 8.1. For $\eta > 0$ small enough, it follows from the results in [EES3] that the index of this operator, after disregarding variations corresponding to conformal reparameterizations, is given by

$$|a| - |b_1| - \ldots - |b_m| + \mu(A) - (n+1)\left(\sum_{i=1}^{m+1} d_i + \sum_{i=1}^{m+1}\sum_{j=1}^{d_i} l^i_j\right).$$

It follows that the linearized $\overline{\partial}$-operator on $\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$ has index

$$I := |a| - |b_1| - \ldots - |b_m| + \mu(A) + \sum_{i=1}^{m+1} d_i - (n+1)\sum_{i=1}^{m+1}\sum_{j=1}^{d_i} l^i_j,$$

after disregarding variations corresponding to conformal reparameterizations, since the dimension of the base is $(n+2)\sum_{i=1}^{m+1} d_i$.

Observe that the fiber F of the bundle

$$\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$$

satisfies

$$\dim F = (n+1)\sum_{i=1}^{m+1}\sum_{j=1}^{d_i} l^i_j.$$

Since

$$T_{(g,c,(\theta^i_j),(s^i_j),(g^i_j))}\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \simeq T_{(g,c,(\theta^i_j),(s^i_j))}\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \oplus F,$$

and since Lemma 8.1 implies that $G = \overline{\partial}$ when restricted to the zero-section $\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \subset \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$, it follows that the linearization of the Fredholm map

$$G = \overline{\partial} \circ \Psi \colon \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$$

has index

$$I + \dim F \quad = \quad |a| - |b_1| - \ldots - |b_m| + \mu(A) + \sum_{i=1}^{m+1} d_i.$$

Recall that the codimension of $Z_{\mathbf{l},\mathbf{v}}$ inside $J^{\mathbf{l}}$ is

$$(n-k)\sum_{i=1}^{m+1} d_i + (n-k-1)\sum_{i=1}^{m+1}\sum_{j=1}^{d_i} l_j^i.$$

The expected dimension of $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ is thus

$$\begin{aligned} I + \dim F - (n-k)\sum_{i=1}^{m+1} d_i - (n-k-1)\sum_{i=1}^{m+1}\sum_{j=1}^{d_i} l_j^i & \\ = \quad |a| - |b_1| - \ldots - |b_m| + \mu(A) - (n-k-1)\left(\sum_{i=1}^{m+1} d_i + \sum_{i=1}^{m+1}\sum_{j=1}^{d_i} l_i^j\right) & \\ = \quad |a| - |b_1| - \ldots - |b_m| + \mu(A) - (n-k-1)\sum_{i=1}^{m+1}\sum_{j=1}^{d_i}(l_i^j + 1) & \\ = \quad |a| - |b_1| - \ldots - |b_m| + \mu(A) - (n-k-1)\sum_{i=1}^{m+1} w_i. & \end{aligned}$$

Similarly, since the codimension of $\widetilde{Z}_{\mathbf{l},\mathbf{v}}$ is 1 inside $Z_{\mathbf{l},\mathbf{v}}$, we conclude that the expected dimension of $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})^{sing}$ is given by

$$|a| - |b_1| - \ldots - |b_m| + \mu(A) - (n-k-1)\sum_{i=1}^{m+1} w_i - 1.$$

8.9. **Transversality and compactness.** Observe that even in the cases when the moduli spaces $\mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$ all are smooth, more work is needed to get a smooth structure on the full moduli spaces $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ in the case when the dimension is greater than zero. Namely, in this case the different spaces $\mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$ have to be glued together along the codimension-1 strata $\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$.

We will start by showing that the moduli spaces $\mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$ can be made smooth by perturbing the almost complex structure J chosen in Section 8.4.

Proposition 8.4. *Any almost complex structure J on $\mathbb{R}\times Y$ satisfying the properties in Section 8.4 can be perturbed outside of a neighborhood of S so that the corresponding moduli spaces*

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$$

and

$$\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$$

become transversely cut out manifolds of the expected dimensions. We call such a choice of J regular*.*

Proof of Proposition 8.4. Let $\widetilde{\mathcal{J}}$ denote the space of almost complex structures on $\mathbb{R} \times Y$ which coincide with some fixed almost complex structure in a neighborhood of $S \subset \mathbb{R} \times Y$ and, moreover, satisfy the properties stated in Section 8.4 in this neighborhood.

Along the lines of the transversality proof in [EES3], we define the configuration spaces

$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) := \bigcup_{J \in \widetilde{\mathcal{J}}} \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \widetilde{\mathcal{J}},$$

$$\mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) := \bigcup_{J \in \widetilde{\mathcal{J}}} \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \widetilde{\mathcal{J}},$$

$$\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) := \bigcup_{J \in \widetilde{\mathcal{J}}} \mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to \widetilde{\mathcal{J}},$$

which are fiber-bundles over the base $\widetilde{\mathcal{J}}$. Furthermore, the maps Ψ and $ev^{\mathbf{l}}$ lift to maps

$$\Psi\colon\ \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) \to \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S),$$

$$ev^{\mathbf{l}}\colon\ \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) \to J^{\mathbf{l}},$$

and the $\overline{\partial}$-operator lifts to a section

$$\overline{\partial}\colon\ \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) \to \mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S)$$

of the fiber-bundle

$$\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) \to \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S).$$

By the results in [EES3] and [Dra] one deduces that the above $\overline{\partial}$-operator is submersive. Furthermore, the variations of the almost complex structure needed for submersivity may be taken to have support in a neighborhood of $[N,+\infty) \times a$, for $N \gg 0$, where the Reeb chord a is the positive asymptotic of the disk. Recall that, by assumption, we can choose $N \gg 0$ such that $[N,+\infty) \times a$ is disjoint from some neighborhood of $S \subset \mathbb{R} \times Y$.

Without loss of generality we can assume that the almost complex structures under consideration are already regular for the moduli spaces $\mathcal{M}_{a;\mathbf{b};A}(V)$, and hence that Proposition 8.3 applies for $J \in \widetilde{\mathcal{J}}$.

Consider a solution

$$u := (g, c, (\theta^i_j), (s^i_j), (g^i_j), J) \in \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S)$$

to the equation

$$G(u) := \overline{\partial} \circ \Psi(u) = 0.$$

First of all, the map

$$ev^{\mathbf{l}}\colon\ \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S) \to J^{\mathbf{l}}$$

is obviously a submersion. Furthermore, the linearization $D_u\overline{\partial}$ is surjective when restricted to variations along the fiber of

$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S)\to(\partial\dot{D})^d_{\mathbf{l}}\times(U_S)^d$$

while keeping the conformal structure fixed (see the proof of Lemma 4.5 in [EES3]). Lemma 8.1 and the form of Ψ near $G^{-1}(0)$ specified by the proof of Proposition 8.3 now imply that

$$D_u(G,ev^{\mathbf{l}})$$

surjective as well.

Using the Sard-Smale theorem, we conclude that there is a Baire subset of compatible almost complex structures $J\in\widetilde{\mathcal{J}}$ for which

$$(G,ev^{\mathbf{l}})\colon\ \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)\to\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)\times J^{\mathbf{l}}$$

is transverse to the two subvarieties

$$\{0\}\times Z_{\mathbf{l},\mathbf{v}}\subset\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)\times J^{\mathbf{l}},$$
$$\{0\}\times\widetilde{Z}_{\mathbf{l},\mathbf{v}}\subset\mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)\times J^{\mathbf{l}}.$$

□

We now prove some transversality results for the full moduli spaces in the rigid case.

Proposition 8.5. *Assume that*

$$\dim\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})=0.$$

For any regular choice of almost complex structure we have

$$\emptyset=\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})\subset\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}).$$

That is, a space $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ *of expected dimension 0 consists of disks having no tangency to* S*.*

Finally, the natural map

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})\to\mathcal{M}_{a;\mathbf{b};A}(V)$$

is injective, and its image is finite.

Proposition 8.6. *Let* $V=\mathbb{R}\times L$*,* $S=\mathbb{R}\times T$*, and let* $\mathbf{v}$ *be an* $\mathbb{R}$*-invariant vector field. Assume that*

$$\dim\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times T,\mathbf{v})=1.$$

For any regular almost complex structure we have

$$\emptyset=\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times T,\mathbf{v})\subset\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times T,\mathbf{v}).$$

That is, a space $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times T,\mathbf{v})$ *of expected dimension 1 consists of disks having no tangency to* $\mathbb{R}\times T$*.*

Finally, the natural map

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times S,\mathbf{v})/\mathbb{R}\to\mathcal{M}_{a;\mathbf{b};A}(\mathbb{R}\times L)/\mathbb{R}$$

is injective, and its image is finite.

Proof of Proposition 8.5 and 8.6. We show Proposition 8.5. The $\mathbb{R}$-invariant case, i.e. Proposition 8.6, follows in the same manner. To that end, observe that in the $\mathbb{R}$-invariant case, the variations of the complex structure in 8.4 may all be taken to be cylindrical.

The statement

$$\emptyset = \mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$$

follows immediately from Proposition 8.4, since

$$\dim \mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) < \dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) = 0.$$

To see that the map

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \to \mathcal{M}_{a;\mathbf{b};A}(V).$$

is an injection, we argue as follows. Suppose that two different configurations in $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ arise from the same pseudo-holomorphic map $g : \dot{D} \to \mathbb{R} \times Y$. Then, g can be interpreted as a solution in

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w}';A}(V,S,\mathbf{v})$$

for some $\mathbf{w}'$ satisfying

$$w'_1 + \ldots + w'_{m+1} > w_1 + \ldots + w_{m+1}.$$

But since $k+1 < n$, the dimension-formula immediately gives

$$\dim \mathcal{M}_{a;\mathbf{b},\mathbf{w}';A}(V,S,\mathbf{v}) < \dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) = 0,$$

which contradicts Proposition 8.4.

It remains to show the compactness of

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V).$$

Since $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ is transversely cut and 0-dimensional and since

$$\mathcal{M}^{sing}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) = \emptyset,$$

it is a discrete subset of $\mathcal{M}_{a;\mathbf{b};A}(V)$.

If $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V)$ would not be compact, this set would have a limit point in

$$\partial\overline{\mathcal{M}}_{a;\mathbf{b};A}(V)$$

which can be identified with a non-trivial holomorphic building. Since the dimension formula is additive, all layers must be rigid with the boundary-point constraints. However, a rigid solution must be a trivial strip over a Reeb chord, which leads to a contradiction. □

9. THE MODULI SPACE OF PSEUDO-HOLOMORPHIC DISKS WITH POINT CONSTRAINTS ON PARALLEL COPIES

As was mentioned in the previous section, obtaining a smooth structure on the moduli spaces $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ when the expected dimension is >0 is a delicate task in general. We will avoid these difficulties by instead constructing spaces consisting of disks with boundary-point constraints on parallel copies of S. These space should be thought of as perturbations of the spaces $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$.

More precisely, for $\delta>0$ and a tuple

$$\mathbf{k}=(k_1,\ldots,k_{w_1+\ldots+w_{m+1}})\in(\mathbb{Z}_{\geq 0})^{w_1+\ldots+w_{m+1}}$$

with increasing indices, we will construct the space

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$$

consisting of pseudo-holomorphic disks with boundary on V and boundary-point constraints on parallel copies of S determined by $\mathbf{v}$, δ and $\mathbf{k}$. Indeed, when the expected dimension is zero, we will show that

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})\simeq\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$$

for all $\delta>0$ small enough.

As in Section 8, we let $V\subset\mathbb{R}\times Y$ be a $(n+1)$-dimensional exact Lagrangian cobordism, $S\subset V$ a $(k+1)$-dimensional submanifold satisfying $k+1<n$ which is cylindrical at the ends, and $\mathbf{v}$ a non-vanishing normal vector-field to S in V which is $\mathbb{R}$-invariant outside of a compact neighborhood. Furthermore, we assume that the Reeb chords on the Legendrian ends of V are disjoint from S outside of a compact set.

Let

$$\begin{cases} d=d_1+\ldots+d_{m+1},\\ \mathbf{w}=(w_1,\ldots,w_{m+1})=(d_1,\ldots,d_{m+1}),\\ \mathbf{l}=((0,\ldots,0),\ldots,(0,\ldots,0))\in\mathbb{Z}^{d_1}\times\ldots\times\mathbb{Z}^{d_{m+1}}.\end{cases}$$

and consider the configuration spaces

$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};A}(V,S;J):=\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)$$

as defined in section 8.1.

Recall that

$$\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J)\subset W^{2,2}_{loc}((\dot{D},\partial\dot{D}),(\mathbb{R}\times Y,V))\times C_{m+1}\times(\partial\dot{D})^d_{\mathbf{l}}\times(U_S)^d,$$

and that there hence is a natural projection

$$ev^0\colon\,\mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};A}(V,S;J)\to(U_S)^d$$

which, moreover, is submersive.

The non-vanishing normal vector-field $\mathbf{v}$ to S induces an identification of a neighborhood of S inside V with a vector bundle

$$N'\oplus\mathbb{R}\to S,$$

under which S corresponds to the zero-section and the $\mathbf{v}$-direction corresponds to the $\mathbb{R}$-summand. Consider the image of the section

$$S_{\delta,i} := (0 \oplus \delta i)(S) \subset N' \oplus \mathbb{R},$$

which corresponds to a "parallel copy" of S. We may furthermore assume that the images of the S_{δ_i} are cylindrical outside of a compact set. We define

$$Z_{\delta,\mathbf{k},\mathbf{v}} := S_{\delta,k_1} \times S_{\delta,k_2} \times \ldots \times S_{\delta,k_d},$$

which we identify with a submanifold $Z_{\delta,\mathbf{k},\mathbf{v}} \subset (U_S)^d$ using the above identification.

Recall that there is a section

$$\overline{\partial}\colon \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};A}(V,S;J) \to \mathcal{E}^{1,\eta}_{a;\mathbf{b},\mathbf{w};0;A}.$$

We now define

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) := (\overline{\partial}, ev^0)^{-1}(\{0\} \times Z_{\delta,\mathbf{k},\mathbf{v}}).$$

By construction, there is an embedding

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial\dot{D})^d_{\mathbf{l}}.$$

By an argument similar to that in Section 8.8, it follows that this spaces has expected dimension

$$\dim \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) = \dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}).$$

For convenience, when $\mathbf{k} = (0,1,\ldots,d-1)$, we will use the notation

$$\mathcal{M}^{\delta}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) := \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}).$$

9.1. **Transversality of the $\overline{\partial}$-operator and compactness.** The following results follow in a manner similar to Proposition 8.4 and Proposition 8.5.

Proposition 9.1. *There is a Baire subset of cylindrical almost complex structures on* $\mathbb{R} \times Y$ *for which*

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$$

is transversely cut-out. Moreover, J *may be arbitrarily chosen outside of a neighborhood of* $\mathbb{R} \times a$.

Proposition 9.2. *Assume that* J *is regular and that*

$$\dim \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) = 0.$$

It follows that the natural map

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \to \mathcal{M}_{a;\mathbf{b};A}(V,S,\mathbf{v})$$

is injective, and that its image is finite.

Furthermore, for any other choice of tuple $\mathbf{k}'$*, there is a cobordism from* $\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ *to* $\mathcal{M}^{\delta,\mathbf{k}'}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$.

Proof. The result follows from a standard transversality argument using Sard's theorem on a smooth finite-dimensional space. ☐

9.2. **A bijection between $\mathcal{M}$ and $\mathcal{M}^{\delta,\mathbf{k}}$ in the rigid case.**

Proposition 9.3. *Suppose that $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ is transversely cut-out and satisfies*

$$\dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) = 0$$

for an almost complex structure J satisfying the properties in Section 8.4 in a neighborhood of S. For any given bound on the indices of $\mathbf{k}$, for all small enough $\delta > 0$, the space $\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ is transversely cut-out, and satisfies

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \simeq \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}).$$

Proposition 9.4. *Let $V = \mathbb{R}\times L$, $S = \mathbb{R}\times T$, and let $\mathbf{v}$ be an $\mathbb{R}$-invariant vector field. Suppose that $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times S,\mathbf{v})$ is transversely cut-out and satisfies*

$$\dim \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times S,\mathbf{v}) = 1$$

for an almost complex structure J satisfying the properties in Section 8.4 in a neighborhood of S. For any given bound on the indices of $\mathbf{k}$, for all small enough $\delta > 0$, the space $\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times S,\mathbf{v})$ is transversely cut-out, and satisfies

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times T,\mathbf{v})/\mathbb{R} \simeq \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(\mathbb{R}\times L,\mathbb{R}\times T,\mathbf{v})/\mathbb{R}.$$

We begin by proving some lemmas.

Lemma 9.5. *Let $f_{\mathbf{v}}(z), \mathbf{v}\in\mathbb{R}^n$ be a smooth family of holomorphic functions in some neighborhood of 0 satisfying*

$$f_{\mathbf{v}}(z) = \sum_{i=0}^{n-1} v_i z^i + \sum_{i=n}^{\infty} a_i(\mathbf{v}) z^i,$$

such that $f_0(z) = z^n h(z)$, where $h(0)\neq 0$. Then, there are neighborhoods $0\in U\subset\mathbb{R}^n$ and $0\in V\subset\mathbb{C}$ such that the elementary symmetric polynomials in n variables evaluated on the n zeros of $f_{\mathbf{v}}(z)$ inside V form a smooth coordinate system on U. In particular, every $f_{\mathbf{v}}$ for $\mathbf{v}\in U$ is uniquely determined by the position of its n zeros inside V.

Proof. We write

$$f_{\mathbf{v}}(z) = p_{\mathbf{v}}(z) + z^{n+1} h_{\mathbf{v}}(z),$$

where $p_{\mathbf{v}}(z) = \sum_{i=0}^{n-1} v_i z^i$ is a polynomial of degree $\leq n$ depending smoothly on $\mathbf{v}$, $h_{\mathbf{v}}(z)$ is holomorphic in z and depends smoothly on $\mathbf{v}$, and where $h_0(z) = h(z)$. We can hence take $\epsilon, \delta > 0$ such that

- $h_{\mathbf{v}}(z)\neq 0$ for $|z|<\epsilon$ and $\|\mathbf{v}\|<\delta$.
- $f_{\mathbf{v}}(z)\neq 0$ for $|z|=\epsilon$ and $\|\mathbf{v}\|<\delta$.

Let $e_i(\mathbf{v})$, $i = 0, \dots, n$ denote the degree-i symmetric polynomial in n variables evaluated on the n zeros $z_1(\mathbf{v}), \dots, z_n(\mathbf{v})$ of $f_{\mathbf{v}}$ inside $D(0, \epsilon)$, where we set $e_0 := 1$. The functions $e_i(\mathbf{v})$ depend smoothly on $\mathbf{v}$ when $\|\mathbf{v}\| < \delta$. This can be seen by induction by first considering the integrals

$$p_k(\mathbf{v}) := \frac{1}{2\pi i} \int_{\partial D(0,\epsilon)} z^k \frac{f'_{\mathbf{v}}(z)}{f_{\mathbf{v}}(z)} dz = \sum_{i=1}^{n} z_i(\mathbf{v})^k, \;\; j = 1, \dots, n,$$

which obviously depend smoothly on $\mathbf{v}$, and then by using Newton's identities

$$k e_k(x_1, \dots, x_n) = \sum_{i=1}^{k} (-1)^{i-1} e_{k-i}(x_1, \dots, x_n) p_i(x_1, \dots, x_n).$$

Furthermore, this equality shows that

$$\left.\frac{\partial}{\partial v_j} e_k\right|_{\mathbf{v}=0} = \left.\frac{\partial}{\partial v_j} \frac{(-1)^{k-1}}{k} p_k\right|_{\mathbf{v}=0},$$

since every factor except e_0 in the terms in Newton's identities vanish for $\mathbf{v} = 0$.

Since $h_{\mathbf{v}}(z)$ has no zero inside $D(0, \epsilon)$ for $\|\mathbf{v}\| < \delta$, and since $f_{\mathbf{v}}(z)$ is nonzero on $\partial D(0, \epsilon)$, for $j, k = 1, \dots, n$ we can calculate

$$\begin{aligned}
\left.\frac{\partial}{\partial v_j} p_k\right|_{\mathbf{v}=0} &= \left.\frac{\partial}{\partial v_j} \frac{1}{2\pi i} \int_{\partial D(0,\epsilon)} z^k \frac{f'_{\mathbf{v}}(z)}{f_{\mathbf{v}}(z)} dz\right|_{\mathbf{v}=0} \\
&= \left.\frac{\partial}{\partial v_j} \frac{1}{2\pi i} \int_{\partial D(0,\epsilon)} z^k \frac{p'_{\mathbf{v}}(z) + (n+1) z^n h_{\mathbf{v}}(z) + z^{n+1} h'_{\mathbf{v}}(z)}{p_{\mathbf{v}}(z) + z^{n+1} h_{\mathbf{v}}(z)} dz\right|_{\mathbf{v}=0} \\
&= \frac{1}{2\pi i} \int_{\partial D(0,\epsilon)} z^k \frac{j z^{j-1} + (n+1) z^n \partial_{v_j} h_{\mathbf{v}}(z)(0) + z^{n+1} \partial_{v_j} h'_{\mathbf{v}}(z)(0)}{z^{n+1} h_0(z)} dz + \\
&\quad - \frac{1}{2\pi i} \int_{\partial D(0,\epsilon)} z^k \frac{((n+1) z^n h_0(z) + z^{n+1} h'_0(z))(z^j + z^{n+1} \partial_{v_j} h_{\mathbf{v}}(z)(0))}{(z^{n+1} h_0(z))^2} dz \\
&= \frac{1}{2\pi i} \int_{\partial D(0,\epsilon)} \frac{j z^{j+k-1}}{z^{n+1} h_0(z)} dz \\
&\quad - \frac{1}{2\pi i} \int_{\partial D(0,\epsilon)} \frac{(z^{j+k+n})((n+1) h_0(z) + z h'_0(z))}{(z^{n+1} h_0(z))^2} dz,
\end{aligned}$$

and after computing the last integral, we arrive at

$$\left.\frac{\partial}{\partial v_j} p_k\right|_{\mathbf{v}=0} = \begin{cases} \frac{i-n-1}{h_0(0)} \neq 0, & j + k = n + 1, \\ -\frac{h'_0(0)}{h_0(0)}, & j + k = n \\ 0, & \text{otherwise.} \end{cases}$$

The lemma now follows by the implicit function theorem. □

Lemma 9.6. *Assume that the almost complex structure is regular. Consider a sequence $\delta_n \to 0$. Every sequence of rigid solutions*

$$g_n \in \mathcal{M}^{\delta_n,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^d_{\mathbf{l}}$$

has a subsequence which converges to a limit $(g,(\varphi_i))$ where

$$g \in \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V).$$

Proof. By the compactness theorem in [BEH+], there is a convergent subsequence inside $\overline{\mathcal{M}}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^d_{\mathbf{l}}$.

For such a sequence, the boundary points which map to $S_{\delta_n,k}$ converge to boundary points mapping to S as $\delta_n \to 0$, and if j such boundary points collide in the limit, this produces a tangency of order j to $\exp_S(\mathbb{R}\mathbf{v})$ of the disk in the limit. Moreover, the tangent of the boundary at that point has a non-negative component in the $\mathbf{v}$-direction in the case when $j > 0$.

Since there are no rigid broken configurations for regular almost complex structures, the statement follows. ☐

Lemma 9.7. *Suppose that $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ satisfies the assumptions of Proposition 9.1. Consider*

$$(g,(\theta^i_j)) \in \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^{d'}_{\mathbf{l}},$$

where g is tangent of order l^i_j to $\exp_S(\mathbb{R}\mathbf{v})$ at θ^i_j, such that the tangent of the boundary at θ^i_j has a positive component in the $\mathbf{v}$-direction when $l^i_j > 0$. For sufficiently small neighborhoods U of $g \in \mathcal{M}_{a;\mathbf{b};A}(V)$ and U^i_j of $\theta^i_j \in \partial \dot{D}$, for each $\delta > 0$ small enough, there is a unique solution

$$(h,(\varphi_i)) \in \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^d_{\mathbf{l}},$$

satisfying $h \in U$, and which has the property that there are exactly $l^i_j + 1$ number of points in $\{\varphi_i\} \cap U^i_j$.

Proof. Suppose that $(g,(\theta^i_j)) \in \mathcal{M}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S,\mathbf{v})$, where $\mathbf{l} = (l^i_j) \in \mathbb{Z}^{d'}$ and $w_1 + \ldots + w_{m+1} = d$.

By Proposition 8.3 there is an embedding

$$\mathcal{M}_{a;\mathbf{b};A}(V)_{U_S} \hookrightarrow \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J),$$

where

$$\mathcal{M}_{a;\mathbf{b};A}(V)_{U_S} \subset \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^{d'}_{\mathbf{l}}$$

is an open submanifold containing $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$. The regularity of J now implies that the map

$$ev^{\mathbf{l}} \colon \mathcal{M}_{a;\mathbf{b};A}(V)_{U_S} \hookrightarrow \mathcal{G}^{2,\eta}_{a;\mathbf{b},\mathbf{w};\mathbf{l};A}(V,S;J) \to J^{\mathbf{l}},$$

induced by restriction is transverse to the varieties

$$Z_{\mathbf{v}},\ \widetilde{Z}_{\mathbf{v}} \subset J^{\mathbf{l}}.$$

Fix a point $(g,(\theta^i_j)) \in \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V,S,\mathbf{v})$ as above, which we identify with a point $(g,(\theta^i_j)) \in \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^{d'}_{\mathbf{l}}$.

We want to construct suitable coordinates around $g \in \mathcal{M}_{a;\mathbf{b};A}(V)$. For convenience we relabel the indices as follows. Let θ_i be the i:th boundary point among the d' points $\{\theta^i_j\} \subset \partial\dot{D}$ relative the order induced by the orientation of $\partial D \setminus \{p_0\}$. Moreover, we let $s_i := g(\theta_i) \in S$ and l_i be the order of tangency of g to $\exp_S(\mathbb{R}\mathbf{v})$ at θ_i, where, by assumption, the tangent of the boundary at θ_i has a positive component in the $\mathbf{v}$-direction when $l_i > 0$.

Since we assume that J satisfies the properties in Section 8.4, we can choose holomorphic coordinates around $s_i \in S$ identifying s_i with the origin in

$$\mathbb{C}^{k+1} \times \mathbb{C} \times \mathbb{C}^{n-k-1},$$

such that, moreover,

- The real-part is identified with V.
- The subset $\mathbb{R}^{k+1} \times \{0\}$ of the real-part is identified with S.
- The vector field $\mathbf{v}$ is identified with the coordinate vector field of the $(k+2)$:th $\mathbb{R}$-factor of the real-part.

We choose holomorphic coordinates on $\partial\dot{D}$ which identifies a neighborhood of $\theta_i \in \dot{D}$ with a neighborhood in the upper half-plane

$$\{(x, y);\ \ y \geq 0\} \subset \mathbb{C},$$

and which identifies θ_i with 0.

For i satisfying $l_i > 0$, Proposition 8.4 implies that we can choose the coordinate x around θ_i so that the component of $g(x)$ projected to the $\mathbb{R} \times \mathbb{R}^{n-k-1}$ factor, i.e. the component of g which is normal to S, is given by

$$(x, x^{l_i+1}\mathbf{h}_i(x))$$

where $\mathbf{h}_i(x)$ is a real-analytic vector-valued function. Also, it follows by Proposition 8.4 that $\mathbf{h}_i(0) \neq 0$, since otherwise this solution would correspond to a solution having negative formal dimension. We may thus assume that the coordinates around s_i chosen above have the property that all components of the vector $\mathbf{h}_i(0)$ are non-zero.

The above coordinates induce locally defined projection maps on the form

$$TJ^{\mathbf{l}} \subset \times_{i=1}^{d'} TJ^{l_i}(\partial\dot{D}, U_S) \to \times_{i=1}^{d'} TJ^{l_i}(\partial\dot{D}, \mathbb{R} \times \mathbb{R}^{n-k-1}).$$

Furthermore, we have the identification

$$TJ^{l_i}(\partial\dot{D}, \mathbb{R} \times \mathbb{R}^{n-k-1}) \simeq \left(TJ^{l_i}(\partial\dot{D}, \mathbb{R}^{n-k-1}) \oplus \mathbb{R} \oplus \mathbb{R}^{l_i}\right) \times \mathbb{R},$$

where the $\mathbb{R}$ and $\mathbb{R}^{l_i}$-summand correspond to the variations of the constant and non-constant part of the germ in the $\mathbf{v}$-direction, respectively. This, in turn, induces a projection

$$\times_{i=1}^{d'} TJ^{l_i}(\partial\dot{D}, \mathbb{R} \times \mathbb{R}^{n-k-1}) \to \times_{i=1}^{d'} \left(TJ^{l_i}(\partial\dot{D}, \mathbb{R}^{n-k-1}) \oplus \mathbb{R}\right).$$

The transversality of $ev^{\mathbf{l}}$ to $Z_{\mathbf{l},\mathbf{v}}$ at $(g,(\theta_1,\ldots,\theta_{d'}))$ implies that $D_{(g,(\theta_1,\ldots,\theta_{d'}))}ev^{\mathbf{l}}$ composed with the projection onto $\bigoplus_{i=1}^{d'}\left(TJ^{l_i}(\partial\dot{D},\mathbb{R}^{n-k-1})\oplus\mathbb{R}\right)$ is an isomorphism.

It follows that there is a section s

$$\begin{tikzcd}\mathcal{M}_{a;\mathbf{b};A}(V) \arrow[r, shift left, "s"] & \mathcal{M}_{a;\mathbf{b};A}(V)\times(\partial\dot{D})^{d'}_{\mathbf{l}} \arrow[l, shift left]\end{tikzcd}$$

defined in a neighborhood of g and satisfying $s(g)=(g,(\theta_1,\ldots,\theta_i))$ such that the composition of $ev^{\mathbf{l}}\circ s$ with the projection

$$\times_{i=1}^{d'}J^{l_i}(\partial\dot{D},\mathbb{R}^{k+1}\times\mathbb{R}\times\mathbb{R}^{n-k-1})\to\times_{i=1}^{d'}A^i,$$

is a local diffeomorphism, where

$$A^i:=\begin{cases}\mathbb{R}^{(n-k-1)(l_i+1)}=J^{l_i}(\partial\dot{D},\mathbb{R}^{n-k-1}), & l_i>0,\\ \mathbb{R}\times\mathbb{R}^{n-k-2}\times\{0\}\subset\mathbb{R}\times\mathbb{R}^{n-k-1}=J^{l_i}(\partial\dot{D},\mathbb{R}\times\mathbb{R}^{n-k-1}), & l_i=0.\end{cases}$$

In conclusion, using the above coordinates, we have found a smooth parametrization

$$\psi\colon\ \times_{i=1}^{d'}\mathbb{R}^{(n-k-1)(l_i+1)}\simeq\times_{i=1}^{d'}J^{l_i}(\partial\dot{D},\mathbb{R}^{n-k-1})\to\mathcal{M}_{a;\mathbf{b};A}(V),$$

of a neighborhood of $g\in\mathcal{M}_{a;\mathbf{b};A}(V)$ such that g corresponds to the origin

We now study the coordinates $\mathbb{R}^{(n-k-1)(l_i+1)}$ corresponding to i, such that $l_i>0$, more carefully. We fix such an i. In the analytic coordinates around $s_i\in U_S$, and the analytic coordinates around $\theta_i(s\circ\psi)\in\partial\dot{D}$ chosen as above, there are coordinates

$$(\mathbf{v}_1,\ldots,\mathbf{v}_d)\in\times_{i=1}^{d'}A^i\simeq\times_{i=1}^{d'}\mathbb{R}^{(n-k-1)(l_i+1)},\ \ \mathbf{v}_i=(\mathbf{v}_{i,1},\ldots,\mathbf{v}_{i,l_i}),$$

in which the pseudo-holomorphic map $\psi(\mathbf{v}_1,\ldots,\mathbf{v}_{d'})$ has component normal to $S\subset V$ given by

$$\left(h_{i,\psi}(x),\mathbf{v}_{i,0}+\mathbf{v}_{i,1}x+\ldots+\mathbf{v}_{i,l_i}x^{l_i}+x^{l_i+1}\mathbf{h}_{i,\psi}(x)\right).$$

Here, $\mathbf{h}_{i,\psi}(x)$ is a real vector-valued analytic function depending smoothly on the coordinates and satisfying $\mathbf{h}_{i,\psi(0)}(x)=\mathbf{h}_i(x)$, and $h_{i,\psi}(x)$ is a real-analytic function depending smoothly on the coordinates and satisfying $h_{i,\psi(0)}(x)=x$. After the coordinate-change in the domain $\widetilde{x}=h_{i,\psi}(x)$ depending on ψ, the map is expressed as

$$\left(\widetilde{x},\widetilde{\mathbf{v}}_{i,0}+\widetilde{\mathbf{v}}_{i,1}\widetilde{x}+\ldots+\widetilde{\mathbf{v}}_{i,l_i}\widetilde{x}^{l_i}+\widetilde{x}^{l_i+1}\widetilde{\mathbf{h}}_{i,\psi}(\widetilde{x})\right),$$

where again $\widetilde{\mathbf{h}}_{i,\psi}(\widetilde{x})$ is a vector-valued analytic function satisfying $\widetilde{\mathbf{h}}_{i,0}(\widetilde{x})=\mathbf{h}_i(\widetilde{x})$, and where the coordinates $\mathbf{v}$ can be expressed in terms of $\widetilde{\mathbf{v}}$.

By Lemma 9.5, together with the above choice of coordinates corresponding to θ_i for which $l_i=0$, we conclude that there is a neighborhood U of $g\in\mathcal{M}_{a;\mathbf{b};A}(V)$ and neighborhoods U_i of the $\theta_i\in\partial\dot{D}$ such that for any sequence

$$0\leq\delta^1_1<\ldots<\delta^1_{l_1+1}<\ldots<\delta^{d'}_1<\ldots<\delta^{d'}_{l_{d'}+1},$$

where $\delta^{d'}_{l_{d'}+1}$ is small enough, there is a unique $f \in U$ and a unique sequence

$$\varphi^1_1 < \ldots < \varphi^1_{l_1+1} < \ldots < \varphi^{d'}_1 < \ldots < \varphi^{d'}_{l_{d'}+1}$$

for which $\varphi^i_j \in U_i$, and the value of $f(\varphi^i_j)$ in the component $\mathbb{R} \times \mathbb{R}^{n-k-1}$ normal to S is $(\delta^i_j, 0)$. Setting the i:th number in $\{\delta^i_j\}_{(i,j)}$ to δk_i, where the former is endowed with the lexicographic order, we conclude that there is a unique solution from

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v})$$

satisfying the required properties. □

Proof of Proposition 9.3 and 9.4. We prove Proposition 9.3, and note that the $\mathbb{R}$-invariant version, i.e. Proposition 9.4, follows in a similar way.

It follows by Proposition 9.2 that one can identify a rigid solution

$$(h, (\varphi_i)) \in \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^d_{\mathbf{l}}$$

with $h \in \mathcal{M}_{a;\mathbf{b};A}(V)$. Using Lemma 9.7, we can therefore find disjoint neighborhoods of each point in

$$\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V),$$

each which contains exactly one point from

$$\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V)$$

for any $\delta > 0$ small enough.

Assume that for every $\delta > 0$ there are solutions in $\mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v})$ not contained the above neighborhoods. Again, using Lemma 9.6, one can produce a sequence of solutions for $\delta_n \to 0$ which converges to a solution in $\mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v}) \subset \mathcal{M}_{a;\mathbf{b};A}(V)$ outside of these neighborhoods, which leads to a contradiction.

To prove that $(h, \varphi_i) \in \mathcal{M}^{\delta,\mathbf{k}}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v})$ is transversely cut out, where $h(\varphi_i) \in S_{i,k_i}$ for $1 \leq i \leq d$, we consider the restriction

$$ev^0 \colon \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^d_{\mathbf{l}} \subset \mathcal{H}^{2,\eta}_{a;\mathbf{b},\mathbf{w};A}(V, S; J) \to (U_S)^d$$

of the evaluation map ev^0.

By the previous arguments, after choosing $\delta > 0$ small enough, we may assume that h is contained in an arbitrary small neighborhood of a solution $g \in \mathcal{M}_{a;\mathbf{b},\mathbf{w};A}(V, S, \mathbf{v})$, where we assume that $g(\theta_i) \in S$ for $1 \leq i \leq d'$, and that g is tangent of order l_i to $\exp_S(\mathbb{R}\mathbf{v})$ at θ_i.

Similarly to the construction in the proof of Lemma 9.7, we can define coordinates

$$\times_{i=1}^{d'} J^{l_i}(\partial \dot{D}, \mathbb{R} \times \mathbb{R}^{n-k-1}) \simeq \times_{i=1}^{d'} \mathbb{R}^{l_i+1} \times \mathbb{R}^{(n-k-1)(l_i+1)}$$

on this neighborhood of $(g, (\theta_i)) \in \mathcal{M}_{a;\mathbf{b};A}(V) \times (\partial \dot{D})^{d'}_{\mathbf{l}}$ in which $(g, (\theta_i))$ corresponds to the origin.

Observe that for i satisfying $l_i = 0$, it follows by Lemma 9.7 that any neighborhood of θ_i contains a unique point $\varphi_j \in \{\varphi_i\}$ for $\delta > 0$ small enough.

We may suppose that the map ev^0 projected to the normal direction of S of the j:th U_S-component coincides with an $\mathbb{R}\times\mathbb{R}^{n-k-1}$-component of the above coordinates around $(g,(\theta_i))\in\mathcal{M}_{a;\mathbf{b};A}(V)\times(\partial\dot{D})^{d'}_{\mathbf{l}}$. Hence, to show the transversality of ev^0, it now suffices to consider the remaining components of ev^0.

The proof of Lemma 9.7 shows that we may assume that for i satisfying $l_i>0$, there are choices of analytic coordinates centered at $\theta_i\in\partial\dot{D}$ and $g(\theta_i)$ in which the components of g normal to S are on the form

$$\left(x,\mathbf{v}_0+\mathbf{v}_1x+\ldots+\mathbf{v}_{l_i}x^{l_i}+\mathcal{O}(x^{l_i+1})\right),$$

where $\{\mathbf{v}_1\ldots,\mathbf{v}_{l_i}\}$ are a subset of the above coordinates around $(g,(\theta_i))\in\mathcal{M}_{a;\mathbf{b};A}(V)\times(\partial\dot{D})^{d'}_{\mathbf{l}}$. The transversality of ev^0 now follows from the fact that the evaluation map at l_i+1 distinct points

$$(x_1,\ldots,x_{l_i+1},\mathbf{v}_0,\mathbf{v}_1,\ldots,\mathbf{v}_{l_i})\mapsto(P(x_1,\mathbf{v}),\ldots,P(x_{l_i+1},\mathbf{v}))$$

is a submersion for

$$P(x,\mathbf{v})=\left(x,\mathbf{v}_0+\mathbf{v}_1x+\ldots+\mathbf{v}_{l_i}x^{l_i}\right),$$

and since, by Lemma 9.7, there are exactly l_i+1 of the distinct points $\{\varphi_j\}$ contained in an arbitrary small neighborhood of θ_i for small enough $\delta>0$. $\square$

10. EXAMPLES

In the following, we assume that the Maslov class vanishes on the involved Legendrian submanifolds

10.1. **0-surgeries.**

We recall the discussions in Section 3 which imply that, for thin enough handle attachments, a Legendrian ambient surgery just depends on the homotopy class of the Legendrian ambient surgery data. In particular, if L is connected, we may suppose that the attaching 0-sphere $S\subset L$ is located in an arbitrarily small subset of L. Since J-holomorphic disks which are rigid with exactly one point-constraint on S come in pairs for 0-spheres S contained in a small enough neighborhood, it follows that the coefficient of a word containing exactly one letter s vanishes for all boundaries $\partial_{S,N}(c)$ in the DGA $(\mathcal{A}(L,S),\partial_{S,N})$.

Since any augmentation of $(\mathcal{A}(L),\partial)$ pulls back to an augmentation of $(\mathcal{A}(L,S),\partial_{S,N})$, and since

$$(\mathcal{A}(L_S),\partial)\simeq(\mathcal{A}(L,S),\partial_{S,N})$$

by Theorem 1.4, we can compute the linearized contact homology of L_S with respect to the pullback of an augmentation ϵ, which yields

$$HLC(L_S;\epsilon)\simeq HLC(\mathcal{A}(L,S);\epsilon)=HLC(L;\epsilon)\oplus\mathbb{Z}_2s,$$

where we have used the fact that $|s|=n-k-1=n-1\neq0$. We refer to [Che] for the definition of linearized contact homology, and its relation with augmentations.

Furthermore, in the case when all Reeb chords c on L have grading $0 \leq |c| \leq 2n-1$, it follows by degree reasons that the coefficient of a word containing at least two letters s vanish for all boundaries $\partial_{S,N}(c)$ in $(\mathcal{A}(L,S),\partial_{N,S})$. Together with the above observation, this leads to the conclusion

$$HC(L_S) \simeq HC(L) * \langle s \rangle,$$

in this case.

The last requirement is for example satisfied by the standard Legendrian sphere $L_{std} \subset J^1(\mathbb{R}^n)$, which has a representation satisfying

$$|c| = n$$

for its only Reeb chord c. L_{std} can be obtained as the Legendrian lift of the *Whitney sphere*, which is the exact Lagrangian immersion an n-sphere in $\mathbb{C}^n$ with one double point defined by the restriction of the map

$$\mathbb{R}^n \times \mathbb{R} \ni (\mathbf{x}, y) \mapsto (1+iy)\mathbf{x} \in \mathbb{C}^n$$

to the unit sphere $S^n \subset \mathbb{R}^n \times \mathbb{R}$.

10.2. **An $(n-2)$-surgery on the standard sphere.** Consider the isotropic $(n-1)$-disk in $\mathbb{C}^n$ having boundary on the Whitney-sphere, which is given by the map

$$\mathbb{R}^{n-1} \ni \mathbf{x}' \mapsto (\mathbf{x}', 0)$$

restricted to the unit disk $D^{n-1} \subset \mathbb{R}^{n-1}$. After a small perturbation, the disk does not pass through the double-point at the origin. We now consider the respective Legendrian lifts to $J^1(\mathbb{R}^n)$ of the Whitney sphere and the isotropic disk, such that the lift of the isotropic disk has boundary on the standard Legendrian sphere L_{std}.

One can perform a Legendrian ambient surgery along the boundary S of the isotropic disk, yielding $L_{std,S} \simeq S^{n-1} \times S^1$. It can be checked by an explicit calculation in the standard complex structure that $\partial_{S,N}(c) = 0$ holds in $(\mathcal{A}(L_{std},S),\partial_{S,N})$. To that end, observe that in the standard complex structure on $\mathbb{C}^n$ there is exactly one family of embedded disks with boundary on the Whitney sphere and a positive puncture at the double point, whose boundaries moreover induce a foliation of the Whitney sphere by embedded arcs.

By Theorem 1.4 it now follows that

$$HC(L_{std,S}) \simeq HC(L_{std}) * \langle s \rangle \simeq \langle c, s \rangle,$$

where $|c| = n$ and $|s| = 1$.

REFERENCES

[BEE] Frédéric Bourgeois, Tobias Ekholm, and Yakov Eliashberg. Effect of Legendrian surgery. *Geom. Topol.*, 16(1):301–389, 2012.

[BEH$^+$] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki, and E. Zehnder. Compactness results in symplectic field theory. *Geom. Topol.*, 7:799–888, 2003.

[Che] Yuri Chekanov. Differential algebra of Legendrian links. *Invent. Math.*, 150(3):441–483, 2002.

[Dra] Dragomir L. Dragnev. Fredholm theory and transversality for noncompact pseudoholomorphic maps in symplectizations. *Comm. Pure Appl. Math.*, 57(6):726–763, 2004.

[EES1] Tobias Ekholm, John Etnyre, and Michael Sullivan. The contact homology of Legendrian submanifolds in $\mathbf{R}^{2n+1}$. *J. Differential Geom.*, 71(2):177–305, 2005.

[EES2] Tobias Ekholm, John Etnyre, and Michael Sullivan. Non-isotopic Legendrian submanifolds in $\mathbf{R}^{2n+1}$. *J. Differential Geom.*, 71(1):85–128, 2005.

[EES3] Tobias Ekholm, John Etnyre, and Michael Sullivan. Legendrian contact homology in $P \times \mathbf{R}$. *Trans. Amer. Math. Soc.*, 359(7):3301–3335 (electronic), 2007.

[EGH] Y. Eliashberg, A. Givental, and H. Hofer. Introduction to symplectic field theory. *Geom. Funct. Anal.*, (Special Volume, Part II):560–673, 2000. GAFA 2000 (Tel Aviv, 1999).

[EHK] T. Ekholm, K. Honda, and T. Kálmán. Legendrian knots and exact Lagrangian cobordisms. preprint (2012).

[Ekh1] Tobias Ekholm. Morse flow trees and Legendrian contact homology in 1-jet spaces. *Geom. Topol.*, 11:1083–1224, 2007.

[Ekh2] Tobias Ekholm. Rational symplectic field theory over $\mathbf{Z}_2$ for exact Lagrangian cobordisms. *J. Eur. Math. Soc. (JEMS)*, 10(3):641–704, 2008.

[EM] Y. Eliashberg and N. Mishachev. *Introduction to the h-principle*, volume 48 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2002.

[ENS] John B. Etnyre, Lenhard L. Ng, and Joshua M. Sabloff. Invariants of Legendrian knots and coherent orientations. *J. Symplectic Geom.*, 1(2):321–367, 2002.

[ES] T. Ekholm and I. Smith. Exact Lagrangian immersions with a single double point. preprint (2011), available at `http://arxiv.org/abs/1111.5932`.

[FR] Dmitry Fuchs and Dan Rutherford. Generating families and Legendrian contact homology in the standard contact space. *J. Topol.*, 4(1):190–226, 2011.

[Gro] M. Gromov. Pseudoholomorphic curves in symplectic manifolds. *Invent. Math.*, 82(2):307–347, 1985.

[HS] John G. Harper and Michael G. Sullivan. A bordered Legendrian contact algebra. preprint (2012), available at `http://arxiv.org/abs/1204.1962`.

[HWZ] H. Hofer, K. Wysocki, and E. Zehnder. Properties of pseudoholomorphic curves in symplectisations. I. Asymptotics. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 13(3):337–379, 1996.

[McD] Dusa McDuff. *Introduction to symplectic topology*, volume 7 of *IAS/Park City Math. Ser.* Amer. Math. Soc., Providence, RI, 1999.

[PC] P. E. Pushkar′ and Yu. V. Chekanov. Combinatorics of fronts of Legendrian links, and Arnol′d's 4-conjectures. *Uspekhi Mat. Nauk*, 60(1(361)):99–154, 2005.

[Sik] Jean-Claude Sikorav. Some properties of holomorphic curves in almost complex manifolds. In *Holomorphic curves in symplectic geometry*, volume 117 of *Progr. Math.*, pages 165–189. Birkhäuser, Basel, 1994.

[Zeh] Kai Zehmisch. *Singularities and Self-Intersections of Holomorphic Discs*. PhD thesis, Universität Leizig, 2009.

Department of mathematics, Uppsala University, Box 480, 751 06 Uppsala, Sweden

E-mail address: `georgios@math.uu.se`